

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 March 2003 (06.03.2003)

PCT

(10) International Publication Number
WO 03/018022 A1

(51) International Patent Classification⁷: **A61K 31/506**,
A61P 35/00, C07D 413/14, 413/12, 417/14, 417/12

(74) Agents: **ODRE, Steven, M.** et al.; Amgen, Inc., One Amgen Center Drive, M/S 27-4-A, Thousand Oaks, CA 91320-1799 (US).

(21) International Application Number: PCT/US02/26926

(22) International Filing Date: 22 August 2002 (22.08.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/314,285 22 August 2001 (22.08.2001) US
10/225,819 21 August 2002 (21.08.2002) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant: **AMGEN INC.** [US/US]; One Amgen Center Drive, Thousand Oaks, CA 91320-1799 (US).

Published:

— with international search report

(72) Inventors: **BUCHANAN, John, L.**; 143 Babcock Street, Brookline, MA 02446 (US). **CHAFFEE, Stuart**; 24 Bates Street, Cambridge, MA 02140 (US). **HARMANCE, Jean-Christophe**; 57 William Street, Andover, MA 01810 (US). **NOVAK, Perry, M.**; 35 Debbie Lane, Milford, MA 01757 (US). **ZHU, Xiaotian**; 131 Coolidge Avenue #324, Watertown, MA 02472 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 2-AMINO-4-HETEROARYLAMINOPYRIMIDINE DERIVATIVES FOR USE IN THE TREATMENT OF CANCER

(57) Abstract: Compounds of Formula (I) are claimed, wherein R¹ is a heteroaryl ring containing at least one oxygen or sulfur heteroatom and R² is as defined in the claims. These compounds are useful in the treatment of cancer.

WO 03/018022 A1

2-AMINO-4-HETEROARYLAMINOPYRIMIDINE DERIVATIVES FOR USE IN THE TREATMENT OF
CANCER

5

FIELD OF THE INVENTION

This invention is in the field of pharmaceutical
agents and specifically relates to compounds, compositions,
10 uses and methods for treating cancer and related disorders.

BACKGROUND OF THE INVENTION

Phosphoryl transferases are a large family of enzymes
15 that transfer phosphorous-containing groups from one
substrate to another. Kinases are a class of enzymes that
function in the catalysis of phosphoryl transfer. The
protein kinases constitute the largest subfamily of
structurally related phosphoryl transferases and are
20 responsible for the control of a wide variety of signal
transduction processes within the cell. Almost all kinases
contain a similar 250-300 amino acid catalytic domain. The
protein kinases may be categorized into families by the
substrates they phosphorylate (e.g., protein-tyrosine,
25 protein-serine/threonine, etc.). Protein kinase sequence
motifs have been identified that generally correspond to
each of these kinase families. Lipid kinases (e.g. PI3K)
constitute a separate group of kinases with structural
similarity to protein kinases.

30 The "kinase domain" appears in a number of
polypeptides which serve a variety of functions. Such
polypeptides include, for example, transmembrane receptors,
intracellular receptor associated polypeptides, cytoplasmic
located polypeptides, nuclear located polypeptides and
35 subcellular located polypeptides. The activity of protein
kinases can be regulated by a variety of mechanisms. It
must be noted, however, that an individual protein kinase

- 2 -

may be regulated by more than one mechanism. These mechanisms include, for example, autophosphorylation, transphosphorylation by other kinases, protein-protein interactions, protein-lipid interactions, protein-
5 polynucleotide interactions, ligand binding, and post-translational modification.

Protein and lipid kinases regulate many different cell processes including, but not limited to, proliferation, growth, differentiation, metabolism, cell cycle events,
10 apoptosis, motility, transcription, translation and other signaling processes, by adding phosphate groups to targets such as proteins or lipids. Phosphorylation events catalyzed by kinases act as molecular on/off switches that can modulate or regulate the biological function of the target
15 protein. Phosphorylation of target proteins occurs in response to a variety of extracellular signals (hormones, neurotransmitters, growth and differentiation factors, etc.), cell cycle events, environmental or nutritional stresses, etc. Protein and lipid kinases can function in
20 signaling pathways to activate or inactivate, or modulate the activity of (either directly or indirectly) the targets. These targets may include, for example, metabolic enzymes, regulatory proteins, receptors, cytoskeletal proteins, ion channels or pumps, or transcription factors. Uncontrolled
25 signaling due to defective control of protein phosphorylation has been implicated in a number of diseases and disease conditions, including, for example, inflammation, cancer, allergy/asthma, disease and conditions of the immune system, disease and conditions of the central
30 nervous system (CNS), cardiovascular disease, dermatology, and angiogenesis.

Initial interest in protein kinases as pharmacological targets was stimulated by the findings that many viral oncogenes encode structurally modified cellular protein

- 3 -

kinases with constitutive enzyme activity. These findings pointed to the potential involvement of oncogene related protein kinases in human proliferative disorders.

Subsequently, deregulated protein kinase activity, resulting
5 from a variety of more subtle mechanisms, has been implicated in the pathophysiology of a number of important human disorders including, for example, cancer, CNS conditions, and immunologically related diseases. The development of selective protein kinase inhibitors that can
10 block the disease pathologies and/or symptoms resulting from aberrant protein kinase activity has therefore generated much interest.

Protein kinases represent a large family of proteins which play a central role in the regulation of a wide
15 variety of cellular processes, maintaining control over cellular function. A partial list of such kinases includes abl, AKT, bcr-abl, Blk, Brk, Btk, c-kit, c-met, c-src, CDK1, CDK2, CDK3, CDK4, CDK5, CDK6, CDK7, CDK8, CDK9, CDK10, cRaf1, CSF1R, CSK, EGFR, ErbB2, ErbB3, ErbB4, Erk, Fak, fes,
20 FGFR1, FGFR2, FGFR3, FGFR4, FGFR5, Fgr, flt-1, Fps, Frk, Fyn, Hck, IGF-1R, INS-R, Jak, KDR, Lck, Lyn, MEK, p38, PDGFR, PIK, PKC, PYK2, ron, tie, tie2, TRK, Yes, and Zap70. Inhibition of such kinases has become an important therapeutic target.

25 A major feature of malignant cells is the loss of control over one or more cell cycle elements. These elements range from cell surface receptors to the regulators of transcription and translation, including the insulin-like growth factors, insulin growth factor-I (IGF-1) and insulin
30 growth factor-2 (IGF-2). [M.J. Ellis, "The Insulin-Like Growth Factor Network and Breast Cancer", Breast Cancer, Molecular Genetics, Pathogenesis and Therapeutics, Humana Press 1999]. The insulin growth factor system consists of families of ligands, insulin growth factor binding proteins,

- 4 -

and receptors. A major physiological role of the IGF-1 system is the promotion of normal growth and regeneration, and overexpressed IGF-1R can initiate mitogenesis and promote ligand-dependent neoplastic transformation.

- 5 Furthermore, IGF-1R plays an important role in the establishment and maintenance of the malignant phenotype.

IGF-1R exists as a heterodimer, with several disulfide bridges. The tyrosine kinase catalytic site and the ATP binding site are located on the cytoplasmic portion of the
10 beta subunit.

Unlike the epidermal growth factor (EGF) receptor, no mutant oncogenic forms of the IGF-1R have been identified. However, several oncogenes have been demonstrated to affect IGF-1 and IGF-1R expression.

- 15 The correlation between a reduction of IGF-1R expression and resistance to transformation has been seen. Exposure of cells to the mRNA antisense to IGF-1R RNA, prevents soft agar growth of several human tumor cell lines.

- Apoptosis is a ubiquitous physiological process used
20 to eliminate damaged or unwanted cells in multicellular organisms. Disregulation of apoptosis is believed to be involved in the pathogenesis of many human diseases. The failure of apoptotic cell death has been implicated in various cancers, as well as autoimmune disorders.
- 25 Conversely, increased apoptosis is associated with a variety of diseases involving cell loss such as neurodegenerative disorders and AIDS. As such, regulators of apoptosis have become an important therapeutic target. It is now established that a major mode of tumor survival is escape
30 from apoptosis. IGF-1R abrogates progression into apoptosis, both in vivo and in vitro. It has also been shown that a decrease in the level of IGF-1R below wild-type levels causes apoptosis of tumor cells in vivo. The ability of IGF-

- 5 -

1R disruption to cause apoptosis appears to be diminished in normal, non-tumorigenic cells.

WO01/00213, published 4 January 2001, describes substituted pyrimidines as SRC kinase inhibitors.

- 5 WO01/40218, published 7 June 2001, describes arylamine derivatives for use as anti-telomerase agents. WO00/39101, published 6 July 2000, describes substituted pyrimidines as anti-cancer agents. WO01/29009, published 26 April 2001, describes substituted pyrimidines as kinase inhibitors.
- 10 WO00/78731, published 28 December 2000, describes cyano substituted pyrimidines as kinase inhibitors. WO00/53595, published 14 September 2000, describes substituted pyrimidines as kinase inhibitors. WO00/39101, published 6 July 2000, describes amino substituted pyrimidines as kinase
- 15 inhibitors. WO00/59892, published 12 October 2000, describes amino substituted pyrimidines as kinase inhibitors. WO97/19065, published 29 May 1997, describes 2-anilino-pyrimidines as kinase inhibitors. EP379806, published 10 April 1996, describes substituted pyrimidines
- 20 for the treatment of neurological disorders. EP1040831, published 4 October 2000, describes substituted pyrimidines as CRF antagonists. Amino substituted pyrimidines were cited in Chem. Abstr. 112:191083. Amino substituted pyrimidines were cited in Chem. Abstr. 72:1114009.
- 25 WO95/33750, published 14 December 1995, describes substituted pyrimidines as CRF antagonists. WO94/26733, published 24 November 1994, describes pyrimidine derivatives as ligands for dopamine receptors. US patent No. 5,958,935 describes substituted pyrimidines as kinase inhibitors. US
- 30 patent No. 4,983,608, describes pyrrolyl-amino substituted pyrimidines as analgesic agents. US patent No. 5,043,317, describes amino substituted pyrimidines as dyes. US patent No. 5,935,966 describes carboxylate substituted pyrimidines as anti-inflammatories. US patent No. 6,080,858 describes a

- 6 -

process for preparing substituted pyrimidines. WO99/50250, published 7 October 1999, describes amino substituted pyrimidines for the treatment of HIV infection. EP945443, published 29 September 1999, describes amino substituted pyrimidines for the treatment of HIV infection. WO99/31073, published 24 June 1999, describes amide substituted pyrimidines. WO00/27825, published 18 May 2000, describes amino substituted pyrimidines for the treatment of HIV infection. WO01/22938, published 5 April 2001, describes amino substituted pyrimidines for the treatment of HIV infection. WO99/41253, published 19 August 1999, describes amino substituted pyrimidines for the treatment of viral infection. WO01/19825, published 22 March 2001, describes amino substituted pyrimidines as synthetic intermediates. WO01/47921, published 5 July 2001, describes amino substituted pyrimidines as kinase inhibitors. WO01/72745, published 4 October 2001, describes 4-heteroaryl-substituted pyrimidines as inhibitors of CDK's. WO01/72717, published 4 October 2001, describes 4-amino-5-cyanopyrimidines as inhibitors of CDK's. WO01/85700, published 15 November 2001, describes pyrimidines as HIV replication inhibitors. WO02/22601, published 21 March 2002, describes 4-(pyrazol-5-ylamino)pyrimidines as kinase inhibitors. WO02/46184, published describes 4-(4-pyrazolyl)-pyrimidines as kinase inhibitors. WO02/46170, published 13 June 2002, describes 2-anilino-pyrimidines as inhibitors of JNK. WO02/46171, published 13 June 2002, describes 2-anilino-pyrimidines as inhibitors of IKK. WO02/47690, published 20 June 2002, describes 4-arylamino-pyrimidines as kinase inhibitors. WO02/48147, published 20 June 2002, describes pyrimidines as kinase inhibitors. WO02/48148, published 20 June 2002, describes pyrimidines as kinase inhibitors. WO02/50065, published 27 June 2002, describes 2-(5-pyrazolylamino)-pyrimidines as kinase inhibitors. Ghoneim et al., Egypt J.

- 7 -

Pharm. Sci., 28, 117-26 (1987)) describe N,N'-bis(3,5-dimethyl-4-isoxazolyl)-6-methyl-2,4-pyrimidinediamine.

WO02/50066, published 27 June 2002, describes 2-(5-pyrazolylamino)-pyrimidines as kinase inhibitors.


5 WO02/57259, published 25 July 2002, describes 4-(5-pyrazolylamino)-pyrimidines as kinase inhibitors.

WO02/59110, published 1 August 2002, describes amino substituted pyrimidines as inhibitors of VEGFR2.

However, compounds of the current invention have not
10 been described as inhibitors for the treatment of cancer.

DESCRIPTION OF THE INVENTION

A class of compounds useful in treating cancer and is
15 defined by Formula I

**I**

wherein R¹ is heteroaryl containing at least one oxygen or
20 sulfur heteroatom, wherein R¹ is optionally substituted
with 1-4 substituents independently selected from R³;
preferably 5- or 6-membered heteroaryl containing at
least one oxygen or sulfur heteroatom, wherein R¹ is
optionally substituted with 1-4 substituents
25 independently selected from R³;
more preferably 5- or 6-membered heteroaryl selected
from isoxazolyl, isothiazolyl, oxazolyl,
thiazolyl, 1,3,4-oxadiazolyl, 1,2,4-oxadiazolyl,
1,3,4-thiadiazolyl and 1,2,4-thiadiazolyl,
30 wherein R¹ is optionally substituted with 1-2
substituents independently selected from R³;

- 8 -

even more preferably isoxazolyl, isothiazolyl,
oxazolyl, thiazolyl, and , wherein R¹ is
optionally substituted with 1-2 substituents
independently selected from R³; and
5 particularly isoxazolyl;
wherein R² is selected from
H,
C₁₋₁₀-alkyl,
C₂₋₁₀-alkenyl,
10 C₂₋₁₀-alkynyl,
C(O)R⁵,
COOR⁵,
C(O)NR⁵R⁵,
S(O)_nR⁵,
15 C₃₋₁₀-cycloalkyl,
C₄₋₁₀-cycloalkenyl,
aryl optionally substituted with 1-5 substituents
independently selected from R³,
R⁴,
20 C₁₋₁₀-alkyl substituted with 1-3 substituents
independently selected from aryl, R⁷ and R⁴,
C₃₋₁₀-cycloalkyl substituted with 1-3 substituents
independently selected from aryl, R⁷ and R⁴, and
C₂₋₁₀-alkenyl substituted with 1-3 substituents
25 independently selected from aryl, R⁷ and R⁴;
preferably C₁₋₆-alkyl,
C₂₋₆-alkenyl,
C₂₋₆-alkynyl,
C₃₋₆-cycloalkyl,
30 C₄₋₆-cycloalkenyl,
R⁴,
phenyl optionally substituted with 1-4 substituents
independently selected from R³,

- 9 -

- C₁₋₆-alkyl substituted with 1-3 substituents
independently selected from aryl, R⁷ and R⁴,
C₃₋₆-cycloalkyl substituted with 1-3 substituents
independently selected from aryl, R⁷ and R⁴, and
5 C₂₋₆-alkenyl substituted with 1-3 substituents
independently selected from aryl and R⁴,
more preferably R⁴ and phenyl optionally substituted
with 1-4 substituents independently selected from
R³,
10 even more preferably 3,4,5-trimethoxyphenyl, 3,4-
dimethoxyphenyl, 2,5-dimethoxyphenyl, 3,4-
dimethoxy-6-methylphenyl, quinolinyl,
benzimidazolyl, indazolyl, 3-
aminosulfonylphenyl and 4-aminosulfonylphenyl;
15 wherein R³ is independently selected from
H,
C₁₋₁₀-alkyl,
C₂₋₁₀-alkenyl,
C₂₋₁₀-alkynyl,
20 C₃₋₁₀-cycloalkyl,
C₄₋₁₀-cycloalkenyl,
aryl,
R⁴,
halo,
25 SR⁵,
OR⁵,
OC(O)R⁵,
NR⁵R⁵,
NR⁵R⁶,
30 COOR⁵,
NO₂,
CN,
C(O)R⁵,
C(O)C(O)R⁵,

- 10 -

- $C(O)NR^5R^5$,
 $S(O)_nR^5$,
 $S(O)_nNR^5R^5$,
 $NR^5C(O)NR^5R^5$,
5 $NR^5C(O)C(O)R^5$,
 $NR^5C(O)R^5$,
 $NR^5(COOR^5)$,
 $NR^5C(O)R^4$,
 $NR^5S(O)_nNR^5R^5$,
10 $NR^5S(O)_nR^5$,
 $NR^5S(O)_nR^4$,
 $NR^5C(O)C(O)NR^5R^5$,
 $NR^5C(O)C(O)NR^5R^6$,
 C_{1-10} -alkyl substituted with 1-3 substituents
15 independently selected from aryl, R^7 and R^4 ; and
 C_{2-10} -alkenyl substituted with 1-3 substituents
independently selected from aryl, R^7 and R^4 ;
preferably selected from
 C_{1-6} -alkyl,
20 C_{2-6} -alkenyl,
 C_{2-6} -alkynyl,
 C_{3-6} -cycloalkyl,
 C_{4-6} -cycloalkenyl,
phenyl,
25 R^4 ,
halo,
 SR^5 ,
 OR^5 ,
 $OC(O)R^5$,
30 NR^5R^5 ,
 NR^5R^6 ,
 $COOR^5$,
 NO_2 ,
CN,

- 11 -

- $C(O)R^5$,
 $C(O)NR^5R^5$,
 $S(O)_nR^5$,
 $S(O)_nNR^5R^5$,
5 $NR^5C(O)NR^5R^5$,
 $NR^5C(O)R^5$,
 $NR^5(COOR^5)$,
 $NR^5C(O)R^4$,
 $NR^5S(O)_nNR^5R^5$,
10 $NR^5S(O)_nR^5$,
 $NR^5S(O)_nR^4$,
 C_{1-6} -alkyl substituted with 1-3 substituents
independently selected from aryl, R^7 and R^4 ; and
 C_{2-6} -alkenyl substituted with 1-3 substituents
15 independently selected from aryl, R^7 and R^4 ;
more preferably C_{1-4} -alkyl, C_{2-4} -alkenyl, C_{2-4} -alkynyl,
phenyl, R^4 , chloro, fluoro, bromo, CF_3 , C_{1-4} -alkoxy,
phenoxy, heterocycloxy, benzyloxy, C_{1-4} -alkyl-
 $C(O)O-$, amino, alkylamino, phenylamino, carboxy, C_{1-}
20 $_4$ -alkoxycarbonyl, NO_2 , CN, C_{1-4} -alkylcarbonyl,
aminocarbonyl, C_{1-4} -alkylaminocarbonyl, C_{1-4} -
alkylsulfonyl, C_{1-4} -alkylaminosulfonyl, benzyl, C_{1-4} -
alkoxyalkyl, C_{1-4} -aminoalkyl, C_{1-4} -alkylaminoalkyl,
and 5-6- membered heterocyclyl- C_{1-4} -alkyl; and
25 even more preferably methyl, ethyl, propyl, tert-
butyl, isopropyl, phenyl, chloro, fluoro, bromo,
 CF_3 , methoxy, phenoxy, benzyloxy, acetyl, amino,
methylamino, phenylamino, carboxy,
ethoxycarbonyl, NO_2 , CN, methylcarbonyl,
30 aminocarbonyl, methylaminocarbonyl,
methylsulfonyl, methylaminosulfonyl, benzyl,
methoxymethyl, aminomethyl, N,N-
dimethylaminoethyl and furylmethyl;

- 12 -

wherein R⁴ is independently a 5-8 membered monocyclic, 8-12 membered bicyclic, or 11-14 membered tricyclic saturated, partially saturated or unsaturated ring system comprising 1-3 heteroatoms if monocyclic, 1-6 heteroatoms if

5 bicyclic, or 1-9 heteroatoms if tricyclic, said heteroatoms independently selected from O, N, or S, which may be saturated or unsaturated, and wherein 0, 1, 2 or 3 atoms of each ring may be substituted by a substituent independently selected from C₁₋₁₀-alkyl, C₂₋₁₀-alkenyl, C₂₋

10 ₁₀-alkynyl, C₃₋₁₀-cycloalkyl, C₄₋₁₀-cycloalkenyl, halo, haloalkyl, sulfo, oxo, SR⁵, OR⁵, NR⁵R⁵, NR⁵R⁶, NR⁶R⁶, COOR⁵, nitro, cyano, S(O)_nR⁵, S(O)_nNR⁵R⁵, C(O)R⁵ and C(O)NR⁵R⁵; preferably a 5-7 membered monocyclic, or 8-11 membered

15 bicyclic saturated, partially saturated or unsaturated ring system comprising 1-3 heteroatoms if monocyclic, or 1-6 heteroatoms if bicyclic, said heteroatoms independently selected from O, N, or S, which may be saturated or unsaturated, and wherein 0, 1, 2 or 3 atoms of each ring may be substituted by a substituent

20 independently selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, halo, C₁₋₆-haloalkyl, oxo, SR⁵, OR⁵, NR⁵R⁵, COOR⁵, nitro, cyano, S(O)_nR⁵, S(O)_nNR⁵R⁵, C(O)R⁵ and C(O)NR⁵R⁵;

25 more preferably 5-6 membered monocyclic, or 8-10 membered bicyclic saturated, partially saturated or unsaturated ring system comprising 1-3 heteroatoms if monocyclic, 1-6 heteroatoms if bicyclic, said heteroatoms independently selected from O, N, or S, which may be saturated or

30 unsaturated, and wherein 0, 1, 2 or 3 atoms of each ring may be substituted by a substituent independently selected from C₁₋₄-alkyl, halo, C₁₋₆-haloalkyl, oxo, OR⁵, NR⁵R⁵, COOR⁵, nitro, cyano, S(O)_nR⁵, S(O)_nNR⁵R⁵, C(O)R⁵ and C(O)NR⁵R⁵; and

- 13 -

even more preferably quinolyl, isoquinolyl,
indazolyl, imidazolyl, pyrazolyl, pyrrolyl,
indolyl, isoindolyl, purinyl, and
naphthyridinyl, wherein R⁴ is optionally
5 substituted by a substituent independently
selected from methyl, isopropyl, tert-butyl,
fluoro, chloro, -CF₃, oxo, methoxy, phenoxy,
amino, methylamino, phenylamino, carboxy,
ethoxycarbonyl, nitro, cyano, methylcarbonyl,
10 aminocarbonyl, methylaminocarbonyl,
methylsulfonyl and methylaminosulfonyl;
wherein R⁵ is independently selected from H, C₁₋₁₀-alkyl, C₂₋₁₀-
alkenyl, C₂₋₁₀-alkynyl, C₃₋₁₀-cycloalkyl, C₄₋₁₀-
cycloalkenyl, R⁴, aryl optionally substituted
15 with 1-3 substituents independently selected from R³,
C_{1-C10} alkyl substituted with 1-3 substituents
independently selected from aryl, R⁷ and R⁴;
C_{3-C10} cycloalkyl substituted with 1-3 substituents
independently selected from aryl, R⁷ and R⁴; and
20 C_{2-C10} alkenyl substituted with 1-3 substituents
independently selected from aryl, R⁷ and R⁴;
preferably selected from H, C₁₋₆-alkyl, and phenyl
optionally substituted with 1-3 substituents
independently selected from R³;
25 more preferably H, C₁₋₆-alkyl, and
phenyl optionally substituted with 1-3 substituents
independently selected from
C₁₋₄-alkyl, chloro, fluoro, bromo, -CF₃, hydroxy,
C₁₋₄-alkoxy, amino, C₁₋₄-alkylamino, carboxy, C₁₋₄-
30 alkoxycarbonyl, -NO₂, -CN, C₁₋₄-alkylcarbonyl, C₁₋₄-
alkylaminocarbonyl, aminocarbonyl,
aminosulfonyl and acetyl;
wherein R⁶ is selected from -C(O)R⁵, -COOR⁵, -C(O)NR⁵R⁵ and -
S(O)_nR⁵;

- 14 -


wherein R^7 is independently halo, $-CF_3$, $-SR^5$, $-OR^5$, $-OC(O)R^5$, $-NR^5R^5$, $-NR^5R^6$, $-NR^6R^6$, $-COOR^5$, $-NO_2$, $-CN$, $-C(O)R^5$, $-OC(O)NR^5R^5$, $-C(O)NR^5R^5$, $-N(R^5)C(O)R^5$, $-N(R^5)(COOR^5)$ and $-S(O)_nNR^5R^5$; and

5 preferably halo, $-OR^5$, $-NR^5R^5$, $-COOR^5$ and $-CN$;

wherein n is 1 or 2;

and pharmaceutically acceptable salts thereof.

The invention also relates to compounds of Formula II



II

10

wherein R^{10} is selected from phenyl, and 5-10 membered heterocyclyl; wherein R^{10} is optionally substituted with 1-2 substituents selected from R^{11} ;

15 wherein R^{11} is selected from C_1 - C_4 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_3 - C_6 cycloalkyl, C_4 - C_6 cycloalkenyl, phenyl, 5-6 membered heterocyclyl, fluoro, chloro, CF_3 , $-OR^{12}$, $-OC(O)R^{12}$, $-NR^{12}R^{12}$, $-COOR^{12}$, $-C(O)R^{12}$, $-C(O)NR^{12}R^{12}$, $-SO_2R^{12}$, $-SO_2NR^{12}R^{12}$, $-NR^{12}C(O)NR^{12}R^{12}$, $-NR^{12}C(O)R^{12}$, $-NR^{12}(COOR^{12})$, $-NR^{12}SO_2NR^{12}R^{12}$, $-NR^{12}SO_2R^{12}$, $-OC(O)NR^{12}R^{12}$, C_1 - C_3 alkyl substituted with 1-3

20

substituents independently selected from optionally substituted phenyl and optionally substituted 5-6 membered heterocyclyl; and


25 C_2 - C_3 alkenyl substituted with 1-3 substituents independently selected from optionally substituted phenyl and optionally substituted 5-6 membered heterocyclyl;

wherein R^{12} is selected from H, C_{1-6} -alkyl, and

- 15 -

phenyl optionally substituted with 1-3 substituents independently selected from C₁₋₄-alkyl, chloro, fluoro, CF₃, hydroxy, C₁₋₄-alkoxy, amino, C₁₋₄-alkylamino, carboxy, C₁₋₄-alkoxycarbonyl, NO₂, CN, C₁₋₄-alkylcarbonyl, C₁₋₄-alkylaminocarbonyl, aminocarbonyl, aminosulfonyl and acetyl; and pharmaceutically acceptable salts thereof.

The invention also relates to compounds of Formula III



III


wherein R¹⁰ is selected from phenyl, and 5-10 membered heterocyclyl; wherein R¹⁰ is optionally substituted with 1-2 substituents selected from R¹¹;
 wherein R¹¹ is selected from C₁-C₄ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₃-C₆ cycloalkyl, C₄-C₆ cycloalkenyl, phenyl, 5-6 membered heterocyclyl, fluoro, chloro, CF₃, -OR¹², -OC(O)R¹², -NR¹²R¹², -COOR¹², -C(O)R¹², -C(O)NR¹²R¹², -SO₂R¹², -SO₂NR¹²R¹², -NR¹²C(O)NR¹²R¹², -NR¹²C(O)R¹², -NR¹²(COOR¹²), -NR¹²SO₂NR¹²R¹², -NR¹²SO₂R¹², -OC(O)NR¹²R¹², C₁-C₃ alkyl substituted with 1-3 substituents independently selected from optionally substituted phenyl and optionally substituted 5-6 membered heterocyclyl; and C₂-C₃ alkenyl substituted with 1-3 substituents independently selected from optionally substituted phenyl and optionally substituted 5-6 membered heterocyclyl;
 wherein R¹² is selected from H, C₁₋₆-alkyl, and phenyl optionally substituted with 1-3 substituents independently selected from

- 16 -

C₁₋₄-alkyl, chloro, fluoro, CF₃, hydroxy, C₁₋₄-alkoxy, amino, C₁₋₄-alkylamino, carboxy, C₁₋₄-alkoxycarbonyl, NO₂, CN, C₁₋₄-alkylcarbonyl, C₁₋₄-alkylaminocarbonyl, aminocarbonyl, aminosulfonyl and acetyl;

5 and pharmaceutically acceptable salts thereof.

The invention also relates to compounds of Formula I'



10

wherein R¹ is heteroaryl containing at least one oxygen or sulfur heteroatom, wherein R¹ is optionally substituted with 1-4 substituents independently selected from R³;

preferably 5- or 6-membered heteroaryl containing at least one oxygen or sulfur heteroatom, wherein R¹ is optionally substituted with 1-3 substituents independently selected from R³;

15 more preferably isoxazolyl, isothiazolyl, oxazolyl, pyranyl, thiazolyl, thiadiazolyl, oxadiazolyl, furazanyl, furyl, and thienyl;

20 even more preferably isoxazolyl, isothiazolyl, oxazolyl, thiazolyl, furyl, furazanyl and thienyl, wherein R¹ is optionally substituted with 1-2 substituents independently selected from R³;

25 particularly isoxazolyl; and

more particularly 3-isoxazolyl or 5-isoxazolyl; wherein R² is selected from R⁴, and aryl optionally substituted with 1-5 substituents independently selected from R³,

30

preferably R⁴ and aryl optionally substituted with 1-3 substituents independently selected from R³;

- 17 -

more preferably naphthyl, 2,3-dihydro-indolyl, 1,3-benzodioxolyl, indolyl, 1,3-dioxo-isoindolyl, indazolyl, pyridyl, quinolyl, isoquinolyl, benzothiazolyl, 1,2,3-benzotriazolyl, benzimidazolyl, and phenyl; wherein R² is optionally substituted with 1-3 substituents independently selected from R³;

even more preferably 2-naphthyl, 2,3-dihydro-indol-6-yl, 1,3-benzodioxol-5-yl, 5-indolyl, 4-indolyl, 1,3-dioxo-isoindol-5-yl, 5-indazolyl, 6-indazolyl, 3-pyridyl, 3-quinolyl, 6-quinolyl, isoquinolyl, benzothiazol-6-yl, benzothiazol-5-yl, 1,2,3-benzotriazol-5-yl, 6-benzimidazolyl, 5-pyridyl, and phenyl;

wherein R² is optionally substituted with 1-3 substituents independently selected from hydroxy, methoxy, ethoxy, cyano, nitro, chloro, fluoro, bromo, dimethylamino, dimethylaminoethyl, 3-dimethylamino-propoxy, methoxycarbonyl, methylcarbonyl, methylcarbonylamino, CH₃C(O)N(CH₃)-, methyl, ethyl, isopropyl, pyrrolidin-1-ylcarbonylethenyl, pyrrolidin-1-ylcarbonylethyl, pyrrolidin-1-ylpropyl, ethynyl, acetyl, ethoxycarbonylbutyl, carboxybutyl, 2-(1-methyl-piperidin-4-yl)-ethoxy, 2-(4-methyl-piperazin-1-yl)ethoxy, 3-(4-methyl-piperazin-1-yl)-propoxy, 3-(piperidin-1-yl)propoxy, 2-piperidin-1-yl-ethoxy, 2-morpholin-4-yl-ethoxy, pentafluoroethyl, trifluoromethyl, trifluoromethoxy, difluoromethoxy, aminocarbonyl, aminosulfonyl, N,N'-di-propylaminosulfonyl, hydroxypropylaminosulfonyl, (2-thiazolyl)aminosulfonyl, butylaminosulfonyl, methylcarbonylaminosulfonyl, methylsulfonyl, 1-methyl-piperidin-4-ylmethoxy, 1-tert-butoxycarbonyl-

- 18 -

piperazin-4-yl, 4-morpholinyl, 4-methylpiperzin-1-yl, 4-piperazinyl, 4-isopropyl-piperazin-1-yl, and oxazol-5-yl; and

particularly 3,4,5-trimethoxyphenyl, 3,4-
5 dimethoxyphenyl, 2,5-dimethoxyphenyl, 3,4-
dimethoxy-6-methylphenyl, 3-
methylcarbonylaminophenyl, 4-[CH₃C(O)NH-SO₂]-
phenyl, 3-isopropylphenyl, 4-
[CH₃C(O)N(CH₃)]phenyl, 3-methoxy-4-
10 morpholinylphenyl, 3-quinolinyl, 6-quinolinyl,
benzimidazolyl, 5-indazolyl, 6-indazolyl, 3-
chlorophenyl, 3-aminosulfonylphenyl and 4-
aminosulfonylphenyl;

wherein R³ is independently selected from

15 H,
C₁₋₁₀-alkyl,
C₂₋₁₀-alkenyl,
C₂₋₁₀-alkynyl,
C₁₋₁₀-haloalkyl,
20 C₃₋₁₀-cycloalkyl,
C₄₋₁₀-cycloalkenyl,
aryl,
R⁴,
halo,
25 SR⁵,
OR⁵,
OC(O)R⁵,
NR⁵R⁵,
NR⁵R⁶,
30 COOR⁵,
nitro,
cyano,
C(O)R⁵,
C(O)C(O)R⁵,

- 19 -

- $C(O)NR^5R^5$,
 $S(O)_nR^5$,
 $S(O)_nNR^5R^5$,
 $S(O)_nNR^5R^6$,
5 $NR^5C(O)NR^5R^5$,
 $NR^5C(O)C(O)R^5$,
 $NR^5C(O)R^5$,
 NR^5COOR^5 ,
 $NR^5C(O)R^4$,
10 $NR^5S(O)_nNR^5R^5$,
 $NR^5S(O)_nR^5$,
 $NR^5S(O)_nR^4$,
 $NR^5C(O)C(O)NR^5R^5$,
 $NR^5C(O)C(O)NR^5R^6$,
15 C_{1-10} -alkyl substituted with 1-3 substituents
independently selected from aryl, R^7 and R^4 ; and
 C_{2-10} -alkenyl substituted with 1-3 substituents
independently selected from aryl, R^7 and R^4 ;
preferably H, C_{1-4} -alkyl, C_{2-4} -alkenyl, C_{2-4} -alkynyl, phenyl,
20 C_{6-10} -cycloalkyl, R^4 , chloro, fluoro, bromo,
trifluoromethyl, hydroxy, C_{1-4} -alkoxy, C_{1-4} -haloalkoxy,
phenoxy, heterocyclyloxy, benzyloxy, C_{1-4} -
alkylcarbonyloxy, amino, alkylamino, phenylamino,
carboxy, C_{1-4} -alkoxycarbonyl, nitro, cyano, C_{1-4} -
25 alkylcarbonyl, aminocarbonyl, C_{1-4} -alkylaminocarbonyl,
 C_{1-4} -alkylsulfonyl, C_{1-4} -alkylaminosulfonyl, benzyl, C_{1-4} -
alkoxyalkyl, C_{1-4} -aminoalkyl, C_{1-4} -alkylaminoalkyl, and
5-6- membered heterocyclyl- C_{1-4} -alkyl;
more preferably H, halo, C_{1-3} -alkyl, C_{2-3} -alkenyl, C_{2-3} -
30 alkynyl, phenyl, C_{6-10} -cycloalkyl, hydroxy, C_{1-3} -
haloalkoxy, C_{1-3} -alkoxy, $-C(O)-C_{1-3}$ -alkyl, and C_{1-3} -
haloalkyl; and

- 20 -

even more preferably H, hydroxy, iodo, methyl, acetyl, trifluoromethyl, methoxy, cyclohexyl, adamantyl, phenyl and trifluoromethoxy;

wherein R⁴ is independently a 5-8 membered monocyclic, 8-12

5 membered bicyclic, or 11-14 membered tricyclic saturated, partially saturated or unsaturated ring system comprising 1-3 heteroatoms if monocyclic, 1-6 heteroatoms if bicyclic, or 1-9 heteroatoms if tricyclic, said heteroatoms independently selected from O, N, or S, which

10 may be saturated or unsaturated, and wherein 0, 1, 2 or 3 atoms of each ring may be substituted by a substituent independently selected from C₁₋₁₀-alkyl, C₂₋₁₀-alkenyl, C₂₋₁₀-alkynyl, C₃₋₁₀-cycloalkyl, C₄₋₁₀-cycloalkenyl, halo, haloalkyl, sulfo, oxo, SR⁵, OR⁵, NR⁵R⁵, NR⁵R⁶, NR⁶R⁶,

15 COOR⁵, nitro, cyano, S(O)_nR⁵, S(O)_nNR⁵R⁵, C(O)R⁵, C(O)NR⁵R⁵ and 6-membered heteroaryl optionally substituted with 1-3 substituents independently selected from R³;

preferably 2,3-dihydro-indolyl, 1,3-benzodioxolyl, indolyl, 1,3-dioxo-isindolyl, indazolyl, pyridyl,

20 quinolyl, isoquinolyl, benzothiazolyl, 1,2,3-benzotriazolyl, benzimidazolyl, and pyridyl; wherein R⁴ is optionally substituted with hydroxy, C₁₋₃-alkoxy, cyano, nitro, halo, C₁₋₃-alkyl, di-C₁₋₃-alkylamino, di-C₁₋₃-alkylamino-C₁₋₃-alkyl, di-C₁₋₃-alkylamino-C₁₋₃-alkoxy, C₁₋₃-alkylcarbonyl, C₁₋₃-alkoxycarbonyl, C₁₋₃-alkylcarbonylamino, pyrrolidinylcarbonyl-C₂₋₃-alkenyl, pyrrolidinylcarbonyl-C₁₋₃-alkyl, pyrrolidinyl-C₁₋₃-alkyl, C₂₋₃-alkynyl, acetyl, C₁₋₃-alkylcarbonyl-C₁₋₃-alkyl, carboxy-C₁₋₃-alkyl, (piperidinyl)-C₁₋₃-alkoxy, (piperazinyl)-C₁₋₃-alkoxy, 2-morpholinyl-C₁₋₃-alkoxy, C₁₋₃-haloalkyl, C₁₋₃-haloalkoxy, aminocarbonyl, aminosulfonyl, C₁₋₃-alkylaminosulfonyl, hydroxy-C₁₋₃-alkylaminosulfonyl, (thiazolyl)aminosulfonyl, C₁₋₄-alkylaminosulfonyl, C₁₋₃-alkylcarbonylaminosulfonyl, C₁₋

30

- 21 -

₃-alkylsulfonyl, C₁₋₃-alkoxycarbonyl-piperazinyl, morpholinyl, C₁₋₃-alkylpiperziny, piperazinyl, C₁₋₃-alkyl-piperazinyl, and oxazolyl;

more preferably 2,3-dihydro-indol-6-yl, 1,3-benzodioxol-5-yl, 5-indolyl, 4-indolyl, 1,3-dioxo-isoindol-5-yl, 5-indazolyl, 6-indazolyl, 3-pyridyl, 3-quinolyl, 6-quinolyl, isoquinolyl, benzothiazol-6-yl, benzothiazol-5-yl, 1,2,3-benzotriazol-5-yl, 6-benzimidazolyl, and 5-pyridyl;

10 wherein R⁵ is independently selected from H, C₁₋₁₀-alkyl, C₂₋₁₀-alkenyl, C₂₋₁₀-alkynyl, C₃₋₁₀-cycloalkyl, C₄₋₁₀-cycloalkenyl, R⁴, aryl optionally substituted with 1-3 substituents independently selected from R³, C₁-C₁₀ alkyl substituted with 1-3 substituents
15 independently selected from aryl, R⁷ and R⁴; C₃-C₁₀ cycloalkyl substituted with 1-3 substituents independently selected from aryl, R⁷ and R⁴; and C₂-C₁₀ alkenyl substituted with 1-3 substituents independently selected from aryl, R⁷ and R⁴;

20 wherein R⁶ is selected from -C(O)R⁵, -COOR⁵, -C(O)NR⁵R⁵ and -S(O)_nR⁵;

wherein R⁷ is independently halo, -CF₃, -SR⁵, -OR⁵, -OC(O)R⁵, -NR⁵R⁵, -NR⁵R⁶, -NR⁶R⁶, -COOR⁵, -NO₂, -CN, -C(O)R⁵, -OC(O)NR⁵R⁵, -C(O)NR⁵R⁵, -N(R⁵)C(O)R⁵, -N(R⁵)(COOR⁵) and -S(O)_nNR⁵R⁵; and


25 wherein n is 1 or 2; preferably 2;

and pharmaceutically acceptable derivative thereof.

The invention also relates to compounds of Formula II'

30

- 22 -



II'

wherein R^{10} is selected from phenyl, and 5-10 membered heterocyclyl; wherein R^{10} is optionally substituted with 1-2 substituents selected from R^{11} ; preferably 3,4,5-trimethoxyphenyl, 3,4-dimethoxy-6-methylphenyl, 3-methoxy-4-morpholinylphenyl and 5-indazolyl;

wherein R^{11} is selected from C_1 - C_4 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_3 - C_{10} cycloalkyl, C_4 - C_6 cycloalkenyl, phenyl, 5-6 membered heterocyclyl, fluoro, chloro, CF_3 , $-OC(O)R^{12}$, $-NR^{12}R^{12}$, $-COOR^2$, $-C(O)R^{12}$, $-C(O)NR^{12}R^{12}$, $-SO_2R^{12}$, $-SO_2NR^{12}R^{12}$, $-NR^{12}C(O)NR^{12}R^{12}$, $-NR^{12}C(O)R^2$, $-NR^{12}(COOR^{12})$, $-NR^{12}SO_2NR^{12}R^{12}$, $-NR^{12}SO_2R^{12}$, $-OC(O)NR^{12}R^{12}$, $-OR^{13}$, C_1 - C_3 alkyl substituted with 1-3 substituents independently selected from optionally substituted phenyl and optionally substituted 5-6 membered heterocyclyl; and C_2 - C_3 alkenyl substituted with 1-3 substituents independently selected from optionally substituted phenyl and optionally substituted 5-6 membered heterocyclyl;

preferably cyclohexyl, adamantyl, phenyl and tert-butyl;

wherein R^{12} is selected from H, C_1 - C_6 -alkyl, and

phenyl optionally substituted with 1-3 substituents


independently selected from

C_1 - C_4 -alkyl, chloro, fluoro, CF_3 , hydroxy, C_1 - C_4 -alkoxy, amino, C_1 - C_4 -alkylamino, carboxy, C_1 - C_4 -alkoxycarbonyl, NO_2 , CN, C_1 - C_4 -alkylcarbonyl, C_1 - C_4 -alkylaminocarbonyl, aminocarbonyl, aminosulfonyl and acetyl; and

wherein R¹³ is selected from H, C₁₋₆-alkyl, amino-C₁₋₆-alkyl, C₁₋₆-alkylamino-C₁₋₆-alkyl, aminocarbonyl-C₁₋₆-alkyl, C₁₋₆-alkylaminocarbonyl-C₁₋₆-alkyl, and phenyl optionally substituted with 1-3 substituents independently

5 selected from
C₁₋₄-alkyl, chloro, fluoro, CF₃, hydroxy, C₁₋₄-alkoxy,
amino, C₁₋₄-alkylamino, carboxy, C₁₋₄-alkoxycarbonyl,
NO₂, CN, C₁₋₄-alkylcarbonyl, C₁₋₄-alkylaminocarbonyl,
aminocarbonyl, aminosulfonyl and acetyl;
10 and pharmaceutically acceptable salts thereof.

The invention also relates to compounds of Formula III'



III'

wherein R¹⁰ is selected from phenyl, and 5-10 membered heterocyclyl; wherein R¹⁰ is optionally substituted with 1-2 substituents selected from R¹¹;

20 preferably 3,4,5-trimethoxyphenyl, 3,4-dimethoxy-6-methylphenyl, 3-methoxy-4-morpholinylphenyl and 5-indazolyl;


wherein R¹¹ is selected from C₁-C₄ alkyl, C₂-C₃ alkenyl, C₂-C₃ alkynyl, C₃-C₁₀ cycloalkyl, C₄-C₆ cycloalkenyl, phenyl, 5-6

25 membered heterocyclyl, fluoro, chloro, CF₃, -OC(O)R¹², -NR¹²R¹², -COOR², -C(O)R¹², -C(O)NR¹²R¹², -SO₂R¹², -SO₂NR¹²R¹², -NR¹²C(O)NR¹²R¹², -NR¹²C(O)R², -NR¹²(COOR¹²), -NR¹²SO₂NR¹²R¹², -NR¹²SO₂R¹², -OC(O)NR¹²R¹², -OR¹³, C₁-C₃ alkyl substituted with 1-3 substituents independently selected

- 24 -

from optionally substituted phenyl and optionally substituted 5-6 membered heterocyclyl; and C₂-C₃ alkenyl substituted with 1-3 substituents independently selected from optionally substituted phenyl and optionally substituted 5-6 membered heterocyclyl; preferably cyclohexyl, adamantyl, phenyl and tert-butyl; wherein R¹² is selected from H, C₁₋₆-alkyl, and phenyl optionally substituted with 1-3 substituents independently selected from C₁₋₄-alkyl, chloro, fluoro, CF₃, hydroxy, C₁₋₄-alkoxy, amino, C₁₋₄-alkylamino, carboxy, C₁₋₄-alkoxycarbonyl, NO₂, CN, C₁₋₄-alkylcarbonyl, C₁₋₄-alkylaminocarbonyl, aminocarbonyl, aminosulfonyl and acetyl; and wherein R¹³ is selected from H, C₁₋₆-alkyl, amino-C₁₋₆-alkyl, C₁₋₆-alkylamino-C₁₋₆-alkyl, aminocarbonyl-C₁₋₆-alkyl, C₁₋₆-alkylaminocarbonyl-C₁₋₆-alkyl, and phenyl optionally substituted with 1-3 substituents independently selected from C₁₋₄-alkyl, chloro, fluoro, CF₃, hydroxy, C₁₋₄-alkoxy, amino, C₁₋₄-alkylamino, carboxy, C₁₋₄-alkoxycarbonyl, NO₂, CN, C₁₋₄-alkylcarbonyl, C₁₋₄-alkylaminocarbonyl, aminocarbonyl, aminosulfonyl and acetyl; and pharmaceutically acceptable salts thereof.

The invention also relates to compounds of Formula IV



IV

- 25 -

wherein R^{10} is selected from phenyl, and 5-10 membered heterocyclyl; wherein R^{10} is optionally substituted with 1-2 substituents selected from R^{11} ;
preferably 3,4,5-trimethoxyphenyl, 3,4-dimethoxy-6-methylphenyl, 3-methoxy-4-morpholinylphenyl and 5-indazolyl;

5 wherein R^{11} is selected from C_1 - C_4 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_3 - C_{10} cycloalkyl, C_4 - C_6 cycloalkenyl, phenyl, 5-6 membered heterocyclyl, fluoro, chloro, CF_3 , $-OC(O)R^{12}$, $-NR^{12}R^{12}$, $-COOR^2$, $-C(O)R^{12}$, $-C(O)NR^{12}R^{12}$, $-SO_2R^{12}$, $-SO_2NR^{12}R^{12}$, $-NR^{12}C(O)NR^{12}R^{12}$, $-NR^{12}C(O)R^2$, $-NR^{12}(COOR^{12})$, $-NR^{12}SO_2NR^{12}R^{12}$, $-NR^{12}SO_2R^{12}$, $-OC(O)NR^{12}R^{12}$, $-OR^{13}$, C_1 - C_3 alkyl substituted with 1-3 substituents independently selected

10 from optionally substituted phenyl and optionally substituted 5-6 membered heterocyclyl; and

15 C_2 - C_3 alkenyl substituted with 1-3 substituents independently selected from optionally substituted phenyl and optionally substituted 5-6 membered heterocyclyl;

20 preferably cyclohexyl, adamantyl, phenyl and tert-butyl; wherein R^{12} is selected from H, C_{1-6} -alkyl, and phenyl optionally substituted with 1-3 substituents independently selected from

25 C_{1-4} -alkyl, chloro, fluoro, CF_3 , hydroxy, C_{1-4} -alkoxy, amino, C_{1-4} -alkylamino, carboxy, C_{1-4} -alkoxycarbonyl, NO_2 , CN, C_{1-4} -alkylcarbonyl, C_{1-4} -alkylaminocarbonyl, aminocarbonyl, aminosulfonyl and acetyl; and

wherein R^{13} is selected from H, C_{1-6} -alkyl, amino- C_{1-6} -alkyl, C_{1-6} -alkylamino- C_{1-6} -alkyl, aminocarbonyl- C_{1-6} -alkyl, C_{1-6} -alkylaminocarbonyl- C_{1-6} -alkyl, and phenyl optionally

30 substituted with 1-3 substituents independently selected from

C_{1-4} -alkyl, chloro, fluoro, CF_3 , hydroxy, C_{1-4} -alkoxy, amino, C_{1-4} -alkylamino, carboxy, C_{1-4} -alkoxycarbonyl,

- 26 -

NO₂, CN, C₁₋₄-alkylcarbonyl, C₁₋₄-alkylaminocarbonyl, aminocarbonyl, aminosulfonyl and acetyl; and pharmaceutically acceptable salts thereof.

5 A family of specific compounds of particular interest within Formula I consists of compounds and pharmaceutically-acceptable salts thereof as follows:

10 N⁴-(5-tert-butylisoxazol-3-yl)-N²-(4,5-dimethoxy-2-methyl-phenyl)-pyrimidine-2,4-diamine; and
N⁴-(5-tert-butylisoxazol-3-yl)-N²-(3,4,5-trimethoxy-phenyl)-pyrimidine-2,4-diamine.

15 A family of specific compounds of particular interest within Formula I consists of compounds and pharmaceutically-acceptable salts thereof as follows:

20 N⁴-(5-tert-Butylisoxazol-3-yl)-N²-(3,4,5-trimethoxy-phenyl)-pyrimidine-2,4-diamine;
N⁴-(5-t-Butyl-isoxazol-3-yl)-N²-(4,5-dimethoxy-2-methyl-phenyl)-pyrimidine-2,4-diamine;
N⁴-(5-t-Butyl-isoxazol-3-yl)-N²-(1H-indazol-5-yl)-pyrimidine-2,4-diamine;
25 N²-(3-Methoxy-4-morpholin-4-yl-phenyl)-N⁴-(5-phenyl-isoxazol-3-yl)-pyrimidine-2,4-diamine;
N²-(3,4,5-trimethoxyphenyl)-N⁴-(5-phenylisoxazol-3-yl)-2,4-pyrimidinediamine;
N²-(2-methyl-4,5-dimethoxyphenyl)-N⁴-(5-phenylisoxazol-3-yl)-2,4-pyrimidinediamine;
30 N²-(2-methyl-4,5-dimethoxyphenyl)-N⁴-(5-adamantylisoxazol-3-yl)-2,4-pyrimidinediamine; and
N²-(3,4,5-Trimethoxyphenyl)-N⁴-(3-cyclohexylisoxazol-5-yl)-2,4-pyrimidinediamine.

- 27 -

Indications

Compounds of the present invention would be useful for, but not limited to, the prevention or treatment of
5 cancer and related diseases. The compounds of the invention have kinase inhibitory activity, such as IGF-1R inhibitory activity. The compounds of the invention are useful in therapy as antineoplasia agents.

Compounds of the invention would be useful for the
10 treatment of neoplasia including cancer and metastasis, including, but not limited to: carcinoma such as cancer of the bladder, breast, colon, kidney, liver, lung (including small cell lung cancer), esophagus, gall-bladder, ovary, pancreas, stomach, cervix, thyroid, prostate, and skin
15 (including squamous cell carcinoma); hematopoietic tumors of lymphoid lineage (including leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell-lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma and Burkett's lymphoma); hematopoietic
20 tumors of myeloid lineage (including acute and chronic myelogenous leukemias, myelodysplastic syndrome and promyelocytic leukemia); tumors of mesenchymal origin (including fibrosarcoma and rhabdomyosarcoma, and other sarcomas, e.g. soft tissue and bone); tumors of the central
25 and peripheral nervous system (including astrocytoma, neuroblastoma, glioma and schwannomas); and other tumors (including melanoma, seminoma, teratocarcinoma, osteosarcoma, xenoderoma pigmentosum, keratocanthoma, thyroid follicular cancer and Kaposi's sarcoma).

- 28 -

The compounds of the present invention are also useful in the treatment of cancer related indications such as solid tumors, sarcomas (especially Ewing's sarcoma and osteosarcoma), retinoblastoma, rhabdomyosarcomas, neuroblastoma, hematopoietic malignancies, including leukemia and lymphoma, tumor-induced pleural or pericardial effusions, and malignant ascites.

The compounds of the present invention are also useful for promoting apoptosis.

10 The compounds of this invention may also act as inhibitors of other protein kinases, e.g. ErbB, KDR, CDK-2, LCK, CDK-5, IKK, JNK3, and thus be effective in the treatment of diseases associated with other protein kinases.

15 Besides being useful for human treatment, these compounds are also useful for veterinary treatment of companion animals, exotic animals and farm animals, including mammals, rodents, and the like. More preferred animals include horses, dogs, and cats.

20 As used herein, the compounds of the present invention include the pharmaceutically acceptable derivatives thereof.

Definitions

25 The term "prevention" includes either preventing the onset of disorders altogether or delaying the onset of a preclinically evident stage of disorders in individuals. This includes prophylactic treatment of those at risk of developing a disease, such as a cancer, for example. "Prophylaxis" is another term for prevention.

30 A "pharmaceutically-acceptable derivative " denotes any salt, ester of a compound of this invention, or any other compound which upon administration to a patient is capable of providing (directly or indirectly) a compound of this invention, or a metabolite or residue thereof,
35 characterized by the ability to treat neoplasia.

- 29 -

The phrase "therapeutically-effective" is intended to qualify the amount of each agent, which will achieve the goal of improvement in disorder severity and the frequency of incidence over treatment of each agent by itself, while
5 avoiding adverse side effects typically associated with alternative therapies. For example, effective neoplastic therapeutic agents prolong the survivability of the patient, inhibit the rapidly-proliferating cell growth associated with the neoplasm, or effect a regression of the neoplasm.

10 The term "H" denotes a single hydrogen atom. This radical may be attached, for example, to an oxygen atom to form a hydroxyl radical.

Where the term "alkyl" is used, either alone or within other terms such as "haloalkyl" and "alkylamino", it
15 embraces linear or branched radicals having one to about twelve carbon atoms. More preferred alkyl radicals are "lower alkyl" radicals having one to about six carbon atoms. Examples of such radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl,
20 isoamyl, hexyl and the like. Even more preferred are lower alkyl radicals having one or two carbon atoms. The term "alkylenyl" embraces bridging divalent alkyl radicals such as methylenyl and ethylenyl.

The term "alkenyl" embraces linear or branched
25 radicals having at least one carbon-carbon double bond of two to about twelve carbon atoms. More preferred alkenyl radicals are "lower alkenyl" radicals having two to about six carbon atoms. Most preferred lower alkenyl radicals are radicals having two to about four carbon atoms. Examples of
30 alkenyl radicals include ethenyl, propenyl, allyl, propenyl, butenyl and 4-methylbutenyl. The terms "alkenyl" and "lower alkenyl", embrace radicals having "cis" and "trans" orientations, or alternatively, "E" and "Z" orientations.

- 30 -

The term "alkynyl" denotes linear or branched radicals having two to about twelve carbon atoms. More preferred alkynyl radicals are "lower alkynyl" radicals having two to about six carbon atoms. Most preferred are lower alkynyl radicals having two to about four carbon atoms. Examples of such radicals include propargyl, butynyl, and the like.

The term "halo" means halogens such as fluorine, chlorine, bromine or iodine atoms.

The terms "ring" and "ring system" refer to a ring comprising the delineated number of atoms, said atoms being carbon or, where indicated, a heteroatom such as nitrogen, oxygen or sulfur. The ring itself, as well as any substituents thereon, may be attached at any atom that allows a stable compound to be formed. The term "nonaromatic" ring or ring system refers to the fact that at least one, but not necessarily all, rings in a bicyclic or tricyclic ring system is nonaromatic.

Leaving groups are species that may be detached from a molecule during a reaction and are known in the art. Examples of such groups include, but are not limited to, halogen groups (e.g., I, Br, F, Cl), sulfonate groups (e.g., mesylate, tosylate), sulfide groups (e.g., SCH₃), and the like. Nucleophiles are species that may be attached to a molecule during reaction and are known in the art. Examples of such groups include, but are not limited to, amines, Grignard reagents, anionic species (e.g., alkoxides, amides, carbanions) and the like.

The term "haloalkyl" embraces radicals wherein any one or more of the alkyl carbon atoms is substituted with halo as defined above. Specifically embraced are monohaloalkyl, dihaloalkyl and polyhaloalkyl radicals. A monohaloalkyl radical, for one example, may have either an iodo, bromo, chloro or fluoro atom within the radical. Dihalo and polyhaloalkyl radicals may have two or more of the same halo

- 31 -

atoms or a combination of different halo radicals. "Lower haloalkyl" embraces radicals having 1-6 carbon atoms. Even more preferred are lower haloalkyl radicals having one to three carbon atoms. Examples of haloalkyl radicals include

5 fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl and dichloropropyl. "Perfluoroalkyl" means

10 alkyl radicals having all hydrogen atoms replaced with fluoro atoms. Examples include trifluoromethyl and pentafluoroethyl.

The term "hydroxyalkyl" embraces linear or branched alkyl radicals having one to about ten carbon atoms any one

15 of which may be substituted with one or more hydroxyl radicals. More preferred hydroxyalkyl radicals are "lower hydroxyalkyl" radicals having one to six carbon atoms and one or more hydroxyl radicals. Examples of such radicals include hydroxymethyl, hydroxyethyl, hydroxypropyl,

20 hydroxybutyl and hydroxyhexyl. Even more preferred are lower hydroxyalkyl radicals having one to three carbon atoms.

The term "alkoxy" embrace linear or branched oxy-containing radicals each having alkyl portions of one to about ten carbon atoms. More preferred alkoxy radicals are

25 "lower alkoxy" radicals having one to six carbon atoms. Examples of such radicals include methoxy, ethoxy, propoxy, butoxy and *tert*-butoxy. Even more preferred are lower alkoxy radicals having one to three carbon atoms. Alkoxy radicals may be further substituted with one or more halo atoms, such

30 as fluoro, chloro or bromo, to provide "haloalkoxy" radicals. Even more preferred are lower haloalkoxy radicals having one to three carbon atoms. Examples of such radicals include fluoromethoxy, chloromethoxy, trifluoromethoxy, trifluoroethoxy, fluoroethoxy and fluoropropoxy.

- 32 -

The term "aryl", alone or in combination, means a carbocyclic aromatic system containing one or two rings wherein such rings may be attached together in a fused manner. The term "aryl" embraces aromatic radicals such as phenyl, naphthyl, indenyl, tetrahydronaphthyl, and indanyl. More preferred aryl is phenyl. Said "aryl" group may have 1 to 5 substituents such as lower alkyl, hydroxyl, halo, lower haloalkyl, nitro, cyano, lower alkoxy and lower alkylamino.

The term "heterocyclyl" embraces saturated, partially saturated and unsaturated heteroatom-containing ring-shaped radicals, where the heteroatoms may be selected from nitrogen, sulfur and oxygen. It does not include rings containing -O-O-, -O-S- or -S-S- portions. Said "heterocyclyl" group may have 1 to 3 substituents such as hydroxyl, halo, haloalkyl, cyano, lower alkyl, lower aralkyl, oxo, lower alkoxy, amino and lower alkylamino.

Examples of saturated heterocyclic radicals include saturated 3 to 6-membered heteromonocyclic group containing 1 to 4 nitrogen atoms [e.g. pyrrolidinyl, imidazolidinyl, piperidinyl, pyrrolinyl, piperazinyl]; saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms [e.g. morpholinyl]; saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms [e.g., thiazolidinyl]. Examples of partially saturated heterocyclyl radicals include dihydrothienyl, dihydropyranyl, dihydrofuryl and dihydrothiazolyl.

Examples of unsaturated heterocyclic radicals, also termed "heteroaryl" radicals, include unsaturated 5 to 6 membered heteromonocyclyl group containing 1 to 4 nitrogen atoms, for example, pyrrolyl, imidazolyl, pyrazolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, triazolyl [e.g., 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl]; unsaturated 5- to 6-membered

- 33 -

heteromonocyclic group containing an oxygen atom, for example, pyranyl, 2-furyl, 3-furyl, etc.; unsaturated 5 to 6-membered heteromonocyclic group containing a sulfur atom, for example, 2-thienyl, 3-thienyl, etc.; unsaturated 5- to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, for example, oxazolyl, isoxazolyl, oxadiazolyl [e.g., 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl]; unsaturated 5 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms, for example, thiazolyl, thiadiazolyl [e.g., 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl].

The term also embraces radicals where heterocyclic radicals are fused/condensed with aryl radicals:

unsaturated condensed heterocyclic group containing 1 to 5 nitrogen atoms, for example, indolyl, isoindolyl, indoliziny, benzimidazolyl, quinolyl, isoquinolyl, indazolyl, benzotriazolyl, tetrazolopyridazinyl [e.g., tetrazolo [1,5-b]pyridazinyl]; unsaturated condensed heterocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms [e.g. benzoxazolyl, benzoxadiazolyl]; unsaturated condensed heterocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms [e.g., benzothiazolyl, benzothiadiazolyl]. Preferred heterocyclic radicals include five to ten membered fused or unfused radicals. More preferred examples of heteroaryl radicals include quinolyl, isoquinolyl, imidazolyl, pyridyl, thienyl, thiazolyl, oxazolyl, furyl, and pyrazinyl. Other preferred heteroaryl radicals are 5- or 6-membered heteroaryl, containing one or two heteroatoms selected from sulfur, nitrogen and oxygen, selected from thienyl, furyl, pyrrolyl, indazolyl, pyrazolyl, oxazolyl, triazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, pyridyl, piperidinyl and pyrazinyl.

- 34 -

The term "sulfonyl", whether used alone or linked to other terms such as alkylsulfonyl, denotes respectively divalent radicals $\text{-SO}_2\text{-}$.

The term "alkylsulfonyl" embraces sulfonyl radicals substituted with an alkyl radical. More preferred alkylsulfonyl radicals are "lower alkylsulfonyl" radicals having one to six carbon atoms. Even more preferred are lower alkylsulfonyl radicals having one to three carbon atoms. Examples of such lower alkylsulfonyl radicals include methylsulfonyl, and ethylsulfonyl.

The terms "sulfamyl," "aminosulfonyl" and "sulfonamidyl," denotes a sulfonyl radical substituted with an amine radical, $(\text{-SO}_2\text{NH}_2)$.

The terms "carboxy" or "carboxyl", whether used alone or with other terms, such as "carboxyalkyl", denotes $\text{-CO}_2\text{H}$.

The term "aralkyl" embraces aryl-substituted alkyl radicals. Preferable aralkyl radicals are "lower aralkyl" radicals having aryl radicals attached to alkyl radicals having one to six carbon atoms. Even more preferred are "phenylalkylenyl" attached to alkyl portions having one to three carbon atoms. Examples of such radicals include benzyl, diphenylmethyl and phenylethyl. The aryl in said aralkyl may be additionally substituted with halo, alkyl, alkoxy, haloalkyl and haloalkoxy.

The term "heterocyclylalkylenyl" embraces heterocyclyl-substituted alkyl radicals. Preferable heterocyclyl alkylenyl radicals are "lower heterocyclylalkylenyl" radicals having heterocyclyl radicals attached to alkyl radicals having one to six carbon atoms. More preferred are heterocyclyl- $\text{C}_1\text{-C}_2$ -alkylenyl radicals such as morpholinylmethyl, methylpiperidinylmethyl, methylpiperazinylmethyl, and the like.

The term "carbonyl", whether used alone or with other terms, such as "aminocarbonyl", denotes -(C=O)- .

- 35 -

The term "alkylamino" embraces "N-alkylamino" and "N,N-dialkylamino" where amino groups are substituted with one alkyl radical and with two alkyl radicals, respectively. More preferred alkylamino radicals are "lower alkylamino" radicals having one or two alkyl radicals of one to six carbon atoms, attached to a nitrogen atom. Even more preferred are lower alkylamino radicals having one to three carbon atoms. Suitable alkylamino radicals may be mono or dialkylamino such as N-methylamino, N-ethylamino, N,N-dimethylamino, N,N-diethylamino or the like.

The term "aminoalkyl" embraces linear or branched alkyl radicals having one to about ten carbon atoms any one of which may be substituted with one or more amino radicals. More preferred aminoalkyl radicals are "lower aminoalkyl" radicals having one to six carbon atoms and one or more amino radicals. Examples of such radicals include aminomethyl, aminoethyl, aminopropyl, aminobutyl and aminohexyl. Even more preferred are lower aminoalkyl radicals having one to three carbon atoms.

The term "alkylaminoalkyl" embraces alkyl radicals substituted with alkylamino radicals. More preferred alkylaminoalkyl radicals are "lower alkylaminoalkyl" radicals having alkyl radicals of one to six carbon atoms. Even more preferred are lower alkylaminoalkyl radicals having alkyl radicals of one to three carbon atoms. Suitable alkylaminoalkyl radicals may be mono or dialkyl, such as N-methylaminomethyl, N,N-dimethylaminoethyl, N,N-diethylaminomethyl and the like.

The term "cycloalkyl" includes saturated carbocyclic groups. Preferred cycloalkyl groups include C₃-C₆ rings. More preferred compounds include, cyclopentyl, cyclopropyl, and cyclohexyl.

The term "cycloalkenyl" includes carbocyclic groups have one or more carbon-carbon double bonds. "Cycloalkenyl"

- 36 -

and "cycloalkyldienyl" compounds are included. Preferred cycloalkenyl groups include C₃-C₆ rings. More preferred compounds include, for example, cyclopentenyl, cyclopentadienyl, cyclohexenyl and cycloheptadienyl.

5 The term "aryloxy" embraces optionally substituted aryl radicals, as defined above, attached to an oxygen atom. Examples of such radicals include phenoxy.

 The term "aralkoxy" embraces oxy-containing aralkyl radicals attached through an oxygen atom to other radicals.
10 More preferred aralkoxy radicals are "lower aralkoxy" radicals having optionally substituted phenyl radicals attached to lower alkoxy radical as described above. The aryl portion may be further substituted.

 The term "heteroaryloxy" embraces optionally
15 substituted heteroaryl radicals, as defined above, attached to an oxygen atom.

 The term "heteroarylalkoxy" embraces heteroarylalkyl radicals attached through an oxygen atom. More preferred heteroarylalkoxy radicals are "lower heteroarylalkoxy"
20 radicals having optionally substituted heteroarylalkyl radicals attached to lower alkoxy radical as described above.

 The term "aminocarbonyl" denotes an amide group of the formula -C(=O)NH₂.

25 The term "alkoxycarbonyl" denotes an ester group, where a carbonyl radical is substituted with an alkoxy radical. More preferred are "lower alkoxycarbonyl" having lower alkoxy radicals as described above attached to a carbonyl radical.

30 The term "alkylcarbonyl" denotes carbonyl groups which have been substituted with an alkyl radical. More preferred are C₁-C₆-alkylcarbonyl radicals, such as methylcarbonyl, ethylcarbonyl and propylcarbonyl.

- 37 -

The terms "N-alkylaminocarbonyl" and "N,N-dialkylaminocarbonyl" denote aminocarbonyl radicals substituted with one or two alkyl radicals, respectively. More preferred are "lower alkylaminocarbonyl" having lower
5 alkyl radicals as described above attached to an aminocarbonyl radical.

A "pharmaceutically acceptable prodrug" means any pharmaceutically acceptable salt, ester, salt of an ester, or other derivative of a compound of this invention which,
10 upon administration to a recipient, is capable of providing (directly or indirectly) a compound of this invention. Particularly favored derivatives and prodrugs are those that increase the bioavailability of the compounds of this invention when such compounds are administered to a mammal
15 (e.g., by allowing an orally administered compound to be more readily absorbed into the blood) or which enhance delivery of the parent compound to a biological compartment (e.g., the brain or lymphatic system) relative to the parent species. Preferred prodrugs include derivatives where a
20 group which enhances aqueous solubility or active transport through the gut membrane is appended to the structure of formulas described herein.

The term "comprising" is meant to be open ended, including the indicated component but not excluding other
25 elements.

The phrase "Formula I-IV" includes subformulas such as I' and III'.

The present invention preferably includes compounds that selectively inhibit IGF-1R.

30 The present invention also comprises the use of a compound of the invention, or pharmaceutically acceptable derivative thereof, in the manufacture of a medicament for the treatment either acutely or chronically of an apoptosis mediated disease state, including those described

- 38 -

previously. The compounds of the present invention are useful in the manufacture of an anti-cancer medicament. The compounds of the present invention are also useful in the manufacture of a medicament to attenuate or prevent
5 disorders through inhibition of IGF-1R.

The present invention comprises a pharmaceutical composition comprising a therapeutically-effective amount of a compound of Formulas I-IV in association with a least one pharmaceutically-acceptable carrier, adjuvant or diluent.

10 The present invention also comprises a method of treating apoptosis related disorders, in a subject, the method comprising treating the subject having or susceptible to such disorder with a therapeutically-effective amount of a compound of the present invention.

15

COMBINATIONS

While the compounds of the invention can be administered as the sole active pharmaceutical agent, they
20 can also be used in combination with one or more compounds of the invention or other agents. When administered as a combination, the therapeutic agents can be formulated as separate compositions that are administered at the same time or sequentially at different times, or the therapeutic
25 agents can be given as a single composition.

The phrase "co-therapy" (or "combination-therapy"), in defining use of a compound of the present invention and another pharmaceutical agent, is intended to embrace administration of each agent in a sequential manner in a
30 regimen that will provide beneficial effects of the drug combination, and is intended as well to embrace co-administration of these agents in a substantially simultaneous manner, such as in a single capsule having a fixed ratio of these active agents or in multiple, separate
35 capsules for each agent.

- 39 -

Specifically, the administration of compounds of the present invention may be in conjunction with additional therapies known to those skilled in the art in the prevention or treatment of neoplasia, such as with radiation
5 therapy or with cytostatic or cytotoxic agents.

If formulated as a fixed dose, such combination products employ the compounds of this invention within the accepted dosage ranges. Compounds of Formula I may also be administered sequentially with known anticancer or cytotoxic
10 agents when a combination formulation is inappropriate. The invention is not limited in the sequence of administration; compounds of formula I may be administered either prior to or after administration of the known anticancer or cytotoxic agent.

15 Currently, standard treatment of primary tumors consists of surgical excision followed by either radiation or IV administered chemotherapy. The typical chemotherapy regime consists of either DNA alkylating agents, DNA intercalating agents, CDK inhibitors, or microtubule
20 poisons. The chemotherapy doses used are just below the maximal tolerated dose and therefore dose limiting toxicities typically include, nausea, vomiting, diarrhea, hair loss, neutropenia and the like.

There are large numbers of antineoplastic agents
25 available in commercial use, in clinical evaluation and in pre-clinical development, which would be selected for treatment of neoplasia by combination drug chemotherapy. Such antineoplastic agents fall into several major categories, namely, antibiotic-type agents, alkylating
30 agents, antimetabolite agents, hormonal agents, immunological agents, interferon-type agents and a category of miscellaneous agents.

A first family of antineoplastic agents which may be used in combination with compounds of the present invention

- 40 -

consists of antimetabolite-type/thymidilate synthase inhibitor antineoplastic agents. Suitable antimetabolite antineoplastic agents may be selected from but not limited to the group consisting of 5-FU-fibrinogen, acanthifolic acid, aminothiadiaazole, brequinar sodium, carmofur, Ciba-Geigy CGP-30694, cyclopentyl cytosine, cytarabine phosphate stearate, cytarabine conjugates, Lilly DATHF, Merrel Dow DDFC, dezaguanine, dideoxycytidine, dideoxyguanosine, didox, Yoshitomi DMDC, doxifluridine, Wellcome EHNA, Merck & Co. EX-015, fazarabine, floxuridine, fludarabine phosphate, 5-fluorouracil, N-(2'-furanidyl)-5-fluorouracil, Daiichi Seiyaku FO-152, isopropyl pyrrolizine, Lilly LY-188011, Lilly LY-264618, methobenzaprim, methotrexate, Wellcome MZPES, norspermidine, NCI NSC-127716, NCI NSC-264880, NCI NSC-39661, NCI NSC-612567, Warner-Lambert PALA, pentostatin, piritrexim, plicamycin, Asahi Chemical PL-AC, Takeda TAC-788, thioguanine, tiazofurin, Erbamont TIF, trimetrexate, tyrosine kinase inhibitors, Taiho UFT and uricytin.

A second family of antineoplastic agents which may be used in combination with compounds of the present invention consists of alkylating-type antineoplastic agents. Suitable alkylating-type antineoplastic agents may be selected from but not limited to the group consisting of Shionogi 254-S, aldo-phosphamide analogues, altretamine, anaxirone, Boehringer Mannheim BBR-2207, bestrabucil, budotitane, Wakunaga CA-102, carboplatin, carmustine, Chinoin-139, Chinoin-153, chlorambucil, cisplatin, cyclophosphamide, American Cyanamid CL-286558, Sanofi CY-233, cyplatate, Degussa D-19-384, Sumimoto DACHP(My₂), diphenylspiromustine, diplatinum cytostatic, Erba distamycin derivatives, Chugai DWA-2114R, ITI E09, elmustine, Erbamont FCE-24517, estramustine phosphate sodium, fotemustine, Unimed G-6-M, Chinoin GYKI-17230, hepsul-fam, ifosfamide, ioprolatin, lomustine, mafosfamide, mitolactol, Nippon

- 41 -

Kayaku NK-121, NCI NSC-264395, NCI NSC-342215, oxaliplatin, Upjohn PCNU, prednimustine, Proter PTT-119, ranimustine, semustine, SmithKline SK&F-101772, Yakult Honsha SN-22, spiromustine, Tanabe Seiyaku TA-077, tauromustine,
5 temozolomide, teroxirone, tetraplatin and trimelamol.

A third family of antineoplastic agents which may be used in combination with compounds of the present invention consists of antibiotic-type antineoplastic agents. Suitable antibiotic-type antineoplastic agents may be selected from
10 but not limited to the group consisting of Taiho 4181-A, aclarubicin, actinomycin D, actinoplanone, Erbamont ADR-456, aeroplysinin derivative, Ajinomoto AN-201-II, Ajinomoto AN-3, Nippon Soda anisomycins, anthracycline, azinomycin-A, bisucaberin, Bristol-Myers BL-6859, Bristol-Myers BMY-25067,
15 Bristol-Myers BMY-25551, Bristol-Myers BMY-26605, Bristol-Myers BMY-27557, Bristol-Myers BMY-28438, bleomycin sulfate, bryostatin-1, Taiho C-1027, calicheomycin, chromoximycin, dactinomycin, daunorubicin, Kyowa Hakko DC-102, Kyowa Hakko DC-79, Kyowa Hakko DC-88A, Kyowa Hakko DC89-A1, Kyowa Hakko
20 DC92-B, ditrissarubicin B, Shionogi DOB-41, doxorubicin, doxorubicin-fibrinogen, elsamicin-A, epirubicin, erbstatin, esorubicin, esperamicin-A1, esperamicin-Alb, Erbamont FCE-21954, Fujisawa FK-973, fostriecin, Fujisawa FR-900482, glidobactin, gregatin-A, grincamycin, herbimycin,
25 idarubicin, illudins, kazusamycin, kesarirhodins, Kyowa Hakko KM-5539, Kirin Brewery KRN-8602, Kyowa Hakko KT-5432, Kyowa Hakko KT-5594, Kyowa Hakko KT-6149, American Cyanamid LL-D49194, Meiji Seika ME 2303, menogaril, mitomycin, mitoxantrone, SmithKline M-TAG, neoenactin, Nippon Kayaku
30 NK-313, Nippon Kayaku NKT-01, SRI International NSC-357704, oxalysine, oxaunomycin, peplomycin, pilatin, pirarubicin, porothramycin, pyrindanycin A, Tobishi RA-I, rapamycin, rhizoxin, rodorubicin, sibanomycin, siwenmycin, Sumitomo SM-5887, Snow Brand SN-706, Snow Brand SN-07, sorangicin-A,

- 42 -

sparsomycin, SS Pharmaceutical SS-21020, SS Pharmaceutical SS-7313B, SS Pharmaceutical SS-9816B, steffimycin B, Taiho 4181-2, talisomycin, Takeda TAN-868A, terpentecin, thrazine, tricrozarin A, Upjohn U-73975, Kyowa Hakko UCN-10028A,
5 Fujisawa WF-3405, Yoshitomi Y-25024 and zorubicin.

A fourth family of antineoplastic agents which may be used in combination with compounds of the present invention consists of a miscellaneous family of antineoplastic agents, including tubulin interacting agents, topoisomerase II
10 inhibitors, topoisomerase I inhibitors and hormonal agents, selected from but not limited to the group consisting of α -carotene, α -difluoromethyl-arginine, acitretin, Biotec AD-5, Kyorin AHC-52, alstonine, amonafide, amphethinile, amsacrine, Angiostat, ankinomycin, anti-neoplaston A10,
15 antineoplaston A2, antineoplaston A3, antineoplaston A5, antineoplaston AS2-1, Henkel APD, aphidicolin glycinate, asparaginase, Avarol, baccharin, batracylin, benfluron, benzotript, Ipsen-Beaufour BIM-23015, bisantrene, Bristo-Myers BMY-40481, Vestar boron-10, bromofosfamide, Wellcome
20 BW-502, Wellcome BW-773, caracemide, carmethizole hydrochloride, Ajinomoto CDAF, chlorsulfaquinoxalone, Chemes CHX-2053, Chemex CHX-100, Warner-Lambert CI-921, Warner-Lambert CI-937, Warner-Lambert CI-941, Warner-Lambert CI-958, clanfenur, claviridenone, ICN compound 1259, ICN
25 compound 4711, Contracan, Yakult Honsha CPT-11, crisnatol, curaderm, cytochalasin B, cytarabine, cytocytin, Merz D-609, DABIS maleate, dacarbazine, datelliptinium, didemnin-B, dihaematoporphyrin ether, dihydrolenperone, dinaline, distamycin, Toyo Pharmar DM-341, Toyo Pharmar DM-75, Daiichi
30 Seiyaku DN-9693, docetaxel elliprabin, elliptinium acetate, Tsumura EPMTTC, the epothilones, ergotamine, etoposide, etretinate, fenretinide, Fujisawa FR-57704, gallium nitrate, genkwadaphnin, Chugai GLA-43, Glaxo GR-63178, grifolan NMF-5N, hexadecylphosphocholine, Green Cross HO-221,

- 43 -

homoharringtonine, hydroxyurea, BTG ICRF-187, ilmofosine, isoglutamine, isotretinoin, Otsuka JI-36, Ramot K-477, Otsuak K-76COONa, Kureha Chemical K-AM, MECT Corp KI-8110, American Cyanamid L-623, leukoregulin, lonidamine, Lundbeck
5 LU-23-112, Lilly LY-186641, NCI (US) MAP, marycin, Merrel Dow MDL-27048, Medco MEDR-340, merbarone, merocyanine derivatives, methylanilinoacridine, Molecular Genetics MGI-136, minactivin, mitonafide, mitoquinone mepidamol, motretinide, Zenyaku Kogyo MST-16, N-(retinoyl)amino acids,
10 Nisshin Flour Milling N-021, N-acylated-dehydroalanines, nafazatrom, Taisho NCU-190, nocodazole derivative, Normosang, NCI NSC-145813, NCI NSC-361456, NCI NSC-604782, NCI NSC-95580, ocreotide, Ono ONO-112, oquizarone, Akzo Org-10172, paclitaxel, pancratistatin, pazelliptine, Warner-
15 Lambert PD-111707, Warner-Lambert PD-115934, Warner-Lambert PD-131141, Pierre Fabre PE-1001, ICRT peptide D, piroxantrone, polyhaematoporphyrin, polypreic acid, Efamol porphyrin, probimane, procarbazine, proglumide, Invitron protease nexin I, Tobishi RA-700, razoxane, Sapporo
20 Breweries RBS, restrictin-P, retelliptine, retinoic acid, Rhone-Poulenc RP-49532, Rhone-Poulenc RP-56976, SmithKline SK&F-104864, Sumitomo SM-108, Kuraray SMANCS, SeaPharm SP-10094, spatol, spirocyclopropane derivatives, spirogermanium, Unimed, SS Pharmaceutical SS-554,
25 strypoldinone, Stypoldione, Suntory SUN 0237, Suntory SUN 2071, superoxide dismutase, Toyama T-506, Toyama T-680, taxol, Teijin TEI-0303, teniposide, thaliblastine, Eastman Kodak TJB-29, tocotrienol, topotecan, Topostin, Teijin TT-82, Kyowa Hakko UCN-01, Kyowa Hakko UCN-1028, ukrain,
30 Eastman Kodak USB-006, vinblastine sulfate, vincristine, vindesine, vinestramide, vinorelbine, vintriptol, vinzolidine, withanolides and Yamanouchi YM-534.

Alternatively, the present compounds may also be used in co-therapies with other anti-neoplastic agents, such as

- 44 -

acemannan, aclarubicin, aldesleukin, alemtuzumab,
alitretinoin, altretamine, amifostine, aminolevulinic acid,
amrubicin, amsacrine, anagrelide, anastrozole, ANCER,
ancestim, ARGLABIN, arsenic trioxide, BAM 002 (Novelos),
5 bexarotene, bicalutamide, broxuridine, capecitabine,
celmoleukin, cetorelix, cladribine, clotrimazole,
cytarabine ocfosphate, DA 3030 (Dong-A), daclizumab,
denileukin diftitox, deslorelin, dexrazoxane, dilazep,
docetaxel, docosanol, doxercalciferol, doxifluridine,
10 doxorubicin, bromocriptine, carmustine, cytarabine,
fluorouracil, HIT diclofenac, interferon alfa, daunorubicin,
doxorubicin, tretinoin, edelfosine, edrecolomab,
eflornithine, emitefur, epirubicin, epoetin beta, etoposide
phosphate, exemestane, exisulind, fadrozole, filgrastim,
15 finasteride, fludarabine phosphate, formestane, fotemustine,
gallium nitrate, gemcitabine, gemtuzumab zogamicin,
gimeracil/oteracil/tegafur combination, glycopine,
goserelin, heptaplatin, human chorionic gonadotropin, human
fetal alpha fetoprotein, ibandronic acid, idarubicin,
20 (imiquimod, interferon alfa, interferon alfa, natural,
interferon alfa-2, interferon alfa-2a, interferon alfa-2b,
interferon alfa-N1, interferon alfa-n3, interferon alfacon-
1, interferon alpha, natural, interferon beta, interferon
beta-1a, interferon beta-1b, interferon gamma, natural
25 interferon gamma-1a, interferon gamma-1b, interleukin-1
beta, iobenguane, irinotecan, irsogladine, lanreotide, LC
9018 (Yakult), leflunomide, lenograstim, lentinan sulfate,
letrozole, leukocyte alpha interferon, leuprorelin,
levamisole + fluorouracil, liarozole, lobaplatin,
30 lonidamine, lovastatin, masoprocyl, melarsoprol,
metoclopramide, mifepristone, miltefosine, mirimostim,
mismatched double stranded RNA, mitoguazone, mitolactol,
mitoxantrone, molgramostim, nafarelin, naloxone +
pentazocine, nartograstim, nedaplatin, nilutamide,

- 45 -

noscapine, novel erythropoiesis stimulating protein, NSC
631570 octreotide, oprelvekin, osaterone, oxaliplatin,
paclitaxel, pamidronic acid, pegaspargase, peginterferon
alfa-2b, pentosan polysulfate sodium, pentostatin,
5 picibanil, pirarubicin, rabbit antithymocyte polyclonal
antibody, polyethylene glycol interferon alfa-2a, porfimer
sodium, raloxifene, raltitrexed, rasburicase, rhenium Re 186
etidronate, RII retinamide, rituximab, romurtide, samarium
(153 Sm) lexicidronam, sargramostim, sizofiran, sobuzoxane,
10 sonermin, strontium-89 chloride, suramin, tasonermin,
tazarotene, tegafur, temoporfin, temozolomide, teniposide,
tetrachlorodecaoxide, thalidomide, thymalfasin, thyrotropin
alfa, topotecan, toremifene, tositumomab-iodine 131,
trastuzumab, treosulfan, tretinoin, trilostane,
15 trimetrexate, triptorelin, tumor necrosis factor alpha,
natural, ubenimex, bladder cancer vaccine, Maruyama vaccine,
melanoma lysate vaccine, valrubicin, verteporfin,
vinorelbine, VIRULIZIN, zinostatin stimalamer, or zoledronic
acid; abarelix; AE 941 (Aeterna), ambamustine, antisense
20 oligonucleotide, bcl-2 (Genta), APC 8015 (Dendreon),
cetuximab, decitabine, dexaminoglutethimide, diaziquone, EL
532 (Elan), EM 800 (Endorecherche), eniluracil, etanidazole,
fenretinide, filgrastim SD01 (Amgen), fulvestrant,
galocitabine, gastrin 17 immunogen, HLA-B7 gene therapy
25 (Vical), granulocyte macrophage colony stimulating factor,
histamine dihydrochloride, ibritumomab tiuxetan, ilomastat,
IM 862 (Cytran), interleukin-2, iproxifene, LDI 200
(Milkhaus), leridistim, lintuzumab, CA 125 MAb (Biomira),
cancer MAb (Japan Pharmaceutical Development), HER-2 and Fc
30 MAb (Medarex), idiotypic 105AD7 MAb (CRC Technology),
idiotypic CEA MAb (Trilex), LYM-1-iodine 131 MAb
(Techniclone), polymorphic epithelial mucin-yttrium 90 MAb
(Antisoma), marimastat, menogaril, mitumomab, motexafin
gadolinium, MX 6 (Galderma), nelarabine, nolatrexed, P 30

- 46 -

protein, pegvisomant, pemetrexed, porfiromycin, prinomastat,
RL 0903 (Shire), rubitecan, satraplatin, sodium
phenylacetate, sparfosic acid, SRL 172 (SR Pharma), SU 5416
(SUGEN), SU 6668 (SUGEN), TA 077 (Tanabe),
5 tetrathiomolybdate, thaliblastine, thrombopoietin, tin ethyl
etiopurpurin, tirapazamine, cancer vaccine (Biomira),
melanoma vaccine (New York University), melanoma vaccine
(Sloan Kettering Institute), melanoma oncolysate vaccine
(New York Medical College), viral melanoma cell lysates
10 vaccine (Royal Newcastle Hospital), or valspodar.

The invention relates to inhibitors of enzymes that
catalyze phosphoryl transfer and/or that bind ATP/GTP
nucleotides, compositions comprising the inhibitors, and
methods of using the inhibitors and inhibitor compositions.
15 The inhibitors and compositions comprising them are useful
for treating or modulating disease in which phosphoryl
transferases, including kinases, may be involved, symptoms
of such disease, or the effect of other physiological events
mediated by phosphoryl transferases, including kinases. The
20 invention also provides for methods of making the inhibitor
compounds and methods for treating diseases in which one or
more phosphoryl transferase, including kinase, activities is
involved.

Alternatively, the present compounds may also be used
25 in co-therapies with other anti-neoplastic agents, such as
other kinase inhibitors including p38 inhibitors and CDK
inhibitors, TNF inhibitors, metallomatrix proteases
inhibitors (MMP), EGFR inhibitors such as Iressa, KDR
inhibitors, COX-2 inhibitors including celecoxib, rofecoxib,
30 parecoxib, valdecoxib, and etoricoxib, NSAID's, SOD mimics
or $\alpha_v\beta_3$ inhibitors.

The present invention comprises a process for the
preparation of a compound of Formula I-IV.

- 47 -

Compounds of the present invention can possess, in general, one or more asymmetric carbon atoms and are thus capable of existing in the form of optical isomers as well as in the form of racemic or non-racemic mixtures thereof.

5 The optical isomers can be obtained by resolution of the racemic mixtures according to conventional processes, *e.g.*, by formation of diastereoisomeric salts, by treatment with an optically active acid or base. Examples of appropriate acids are tartaric, diacetyltartaric, dibenzoyltartaric,
10 ditoluoyltartaric, and camphorsulfonic acid and then separation of the mixture of diastereoisomers by crystallization followed by liberation of the optically active bases from these salts. A different process for separation of optical isomers involves the use of a chiral
15 chromatography column optimally chosen to maximize the separation of the enantiomers. Still another available method involves synthesis of covalent diastereoisomeric molecules by reacting compounds of the invention with an optically pure acid in an activated form or an optically
20 pure isocyanate. The synthesized diastereoisomers can be separated by conventional means such as chromatography, distillation, crystallization or sublimation, and then hydrolyzed to deliver the enantiomerically pure compound. The optically active compounds of the invention can likewise
25 be obtained by using active starting materials. These isomers may be in the form of a free acid, a free base, an ester or a salt.

Compounds of the present invention can possess, in general, tautomeric forms, which are included in the family
30 of compounds in Formula I-IV.

Also included in the family of compounds of Formula I-IV are the pharmaceutically-acceptable salts thereof. The term "pharmaceutically-acceptable salts" embraces salts commonly used to form alkali metal salts and to form

- 48 -

addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically-acceptable. Suitable pharmaceutically-acceptable acid addition salts of compounds of Formula I-IV

5 may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, arylaliphatic,

10 heterocyclic, carboxylic and sulfonic classes of organic acids, example of which are acetic, adipic, algenic, anthranilic, ascorbic, aspartic, benzoic, benzenesulfonic, butyric, camphoric, camphorsulfonic, citric, cyclopentanepropionic, cyclohexylaminosulfonic, digluconic,

15 dodecylsulfonic, ethanesulfonic, formic, fumaric, galactaric, galacturonic, glycolic, gluconic, glucuronic, glucoheptanoic, glutamic, glycerophosphonic, heptanoic, hexanoic, 4-hydroxybenzoic, 2-hydroxyethanesulfonic, β -hydroxybutyric, lactic, malic, maleic, mandelic, mesylic,

20 methanesulfonic, nicotinic, 2-naphthalenesulfonic, oxalic, palmoic, pectinic, pivalic, persulfuric, 2-phenylpropionic, picric, pyruvic, propionic, phenylacetic, embonic (pamoic), cyclopentane proprionic, pantothenic, toluenesulfonic, salicylic, sulfanilic, stearic, succinic, tartaric,

25 thiocyanic, and undecanoic.

Suitable pharmaceutically-acceptable base addition salts of compounds of Formula I-IV include metallic salts, such as salts made from alkali metals and alkaline earth metals including, for example, aluminum, calcium, lithium,

30 magnesium, potassium, sodium and zinc, or salts made from organic bases including primary, secondary and tertiary amines, substituted amines including cyclic amines, such as caffeine, arginine, diethylamine, N-ethyl piperidine, histidine, glucamine, isopropylamine, lysine, morpholine, N-

- 49 -

ethyl morpholine, piperazine, piperidine, ammonia, triethylamine, trimethylamine. All of these salts may be prepared by conventional means from the corresponding compound of the invention by reacting, for example, the
5 appropriate acid or base with the compound of Formula I-IV.

Also, the basic nitrogen-containing groups can be quaternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl,
10 and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides, and others. Water or oil-soluble or dispersible products are thereby obtained.

15 Additional examples of such salts can be found in Berge et al., J. Pharm. Sci., 66, 1 (1977).

Combinations of substituents and variables envisioned by this invention are only those that result in the formation of stable compounds. The term "stable", as used
20 herein, refers to compounds which possess stability sufficient to allow preparation.


As used herein, the compounds of this invention, including the compounds described herein, are defined to include pharmaceutically acceptable derivatives or prodrugs
25 thereof.

Pharmaceutically acceptable salts of the compounds of this invention include those derived from pharmaceutically acceptable inorganic and organic acids and bases. Other acids, such as oxalic, while not in themselves
30 pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining the compounds of the invention and their pharmaceutically acceptable acid addition salts. This invention also envisions the quaternization of any basic nitrogen-

- 50 -


containing groups of the compounds disclosed herein. Water or oil-soluble or dispersible products may be obtained by such quaternization.

The invention relates to a process for making a
 5 compound of any of the formulas described herein, comprising reacting a pyrimidine of one or more of the formulas:



10 with an appropriate nucleophilic agent or agents, wherein the groups in said formulas are as defined herein.

The invention also relates to a process for making a
 compound of any of the formulas described herein, comprising
 15 reacting a pyrimidine of one or more of the formulas:




with an appropriate nucleophilic agent or agents, wherein L
 20 is defined as a leaving group and the groups in said formulas are as defined herein.

GENERAL SYNTHETIC PROCEDURES

25 The compounds of the invention can be synthesized according to the following procedures of Schemes 1-2, wherein the substituents are as defined for Formulas I-IV, above, except where further noted.

30


- 51 -

Scheme 1

5 Monoamine substituted pyrimidines 2 can be prepared according to the method set out in Scheme 1. 2,4-Dichloropyrimidine 1 is coupled with heteroarylamines, in the presence of base, such as NaH, and a solvent, such as DMF or THF, at a temperature of about 0 °C to about RT to
 10 give (2-chloro-pyrimidin-4-yl)amine 2.

Alternatively, 2,4-dichloropyrimidine 1 is coupled with an amine in the presence of NaOt-Bu, in a solvent, such as t-BuOH, at a temperature about RT to yield monoamine-substituted pyrimidines 2.

15

Scheme 2

20 Monoamine substituted pyrimidines 2 are reacted with an amine having an active hydrogen, such as R^2NH_2 , in solvent, such as acetone and water, and in the presence of acid, such as conc. HCl, to give the diamine substituted pyrimidine 3.

25 Alternatively, the reaction can be performed in a solvent such as IPA or DMSO, with or without DIEA or in a solvent such as IPA or DMSO with or without $Et_3N \cdot TFA$, or in a solvent such as HOAc.

- 52 -

Preferably the reaction is heated, more preferably at a temperature of about $>50^{\circ}\text{C}$, even more preferably at a temperature of about $90-100^{\circ}\text{C}$.

If one or more other functional groups, for example
5 carboxy, hydroxy, amino, or mercapto, are or need to be
protected in a compound of Formulas I-IV, because they
should not take part in the reaction, these are such groups
as are usually used in the synthesis of peptide compounds,
and also of cephalosporins and penicillins, as well as
10 nucleic acid derivatives and sugars.

The protecting groups may already be present in
precursors and should protect the functional groups
concerned against unwanted secondary reactions, such as
acylations, etherifications, esterifications, oxidations,
15 solvolysis, and similar reactions. It is a characteristic of
protecting groups that they lend themselves readily, i.e.
without undesired secondary reactions, to removal, typically
by solvolysis, reduction, photolysis or also by enzyme
activity, for example under conditions analogous to
20 physiological conditions, and that they are not present in
the end-products. The specialist knows, or can easily
establish, which protecting groups are suitable with the
reactions mentioned above and hereinafter.

The protection of such functional groups by such
25 protecting groups, the protecting groups themselves, and
their removal reactions are described for example in
standard reference works, such as J. F. W. McOmie,
"Protective Groups in Organic Chemistry", Plenum Press,
London and New York 1973, in T. W. Greene, "Protective
30 Groups in Organic Synthesis", Wiley, New York 1981, in "The
Peptides"; Volume 3 (editors: E. Gross and J. Meienhofer),
Academic Press, London and New York 1981, in "Methoden der
organischen Chemie" (Methods of organic chemistry), Houben
Weyl, 4th edition, Volume 15/1, Georg Thieme Verlag,

- 53 -

Stuttgart 1974, in H.-D. Jakubke and H. Jescheit,
"Aminosäuren, Peptide, Proteine" (Amino acids, peptides,
proteins), Verlag Chemie, Weinheim, Deerfield Beach, and
Basel 1982, and in Jochen Lehmann, "Chemie der
5 Kohlenhydrate: Monosaccharide und Derivate" (Chemistry of
carbohydrates: monosaccharides and derivatives), Georg
Thieme Verlag, Stuttgart 1974.

In the additional process steps, carried out as
desired, functional groups of the starting compounds which
10 should not take part in the reaction may be present in
unprotected form or may be protected for example by one or
more of the protecting groups mentioned above under
"protecting groups". The protecting groups are then wholly
or partly removed according to one of the methods described
15 there.

Salts of a compound of formula I with a salt-forming
group may be prepared in a manner known *per se*. Acid
addition salts of compounds of formula I may thus be
obtained by treatment with an acid or with a suitable anion
20 exchange reagent. A salt with two acid molecules (for
example a dihalogenide of a compound of formula I) may also
be converted into a salt with one acid molecule per compound
(for example a monohalogenide); this may be done by heating
to a melt, or for example by heating as a solid under a high
25 vacuum at elevated temperature, for example from 130-170 °C,
one molecule of the acid being expelled per molecule of a
compound of formula I.

Salts can usually be converted to free compounds, e.g.
by treating with suitable basic agents, for example with
30 alkali metal carbonates, alkali metal hydrogen carbonates,
or alkali metal hydroxides, typically potassium carbonate or
sodium hydroxide.

All process steps described here can be carried out
under known reaction conditions, preferably under those

- 54 -

specifically mentioned, in the absence of or usually in the presence of solvents or diluents, preferably such as are inert to the reagents used and able to dissolve these, in the absence or presence of catalysts, condensing agents or
5 neutralizing agents, for example ion exchangers, typically cation exchangers, for example in the H^+ form, depending on the type of reaction and/or reactants at reduced, normal, or elevated temperature, for example in the range from about -100 °C to about 190 °C, preferably from about -80 °C to
10 about 150 °C, for example at about -80 to about 60 °C, at RT, at about -20 to about 40 °C or at the boiling point of the solvent used, under atmospheric pressure or in a closed vessel, where appropriate under pressure, and/or in an inert atmosphere, for example, under argon or nitrogen.

15 Salts may be present in all starting compounds and transients, if these contain salt-forming groups. Salts may also be present during the reaction of such compounds, provided the reaction is not thereby disturbed.

In certain cases, typically in hydrogenation
20 processes, it is possible to achieve stereoselective reactions, allowing for example easier recovery of individual isomers.

The solvents from which those can be selected which are suitable for the reaction in question include, for
25 example, H_2O , esters, typically lower alkyl-lower alkanoates, e.g. EtOAc, ethers, typically aliphatic ethers, e.g. Et_2O , or cyclic ethers, e.g. THF, liquid aromatic hydrocarbons, typically benzene or toluene, alcohols, typically MeOH, EtOH or 1-propanol, IPA, nitriles, typically
30 CH_3CN , halogenated hydrocarbons, typically CH_2Cl_2 , acid amides, typically DMF, bases, typically heterocyclic nitrogen bases, e.g. pyridine, carboxylic acids, typically lower alkanecarboxylic acids, e.g. HOAc, carboxylic acid anhydrides, typically lower alkane acid anhydrides, e.g.

- 55 -

acetic anhydride, cyclic, linear, or branched hydrocarbons, typically cyclohexane, hexane, or isopentane, or mixtures of these solvents, e.g. aqueous solutions, unless otherwise stated in the description of the process.

5 The invention relates also to those forms of the process in which one starts from a compound obtainable at any stage as a transient and carries out the missing steps, or breaks off the process at any stage, or forms a starting material under the reaction conditions, or uses said
10 starting material in the form of a reactive derivative or salt, or produces a compound obtainable by means of the process according to the invention and processes the said compound *in situ*. In the preferred embodiment, one starts from those starting materials which lead to the compounds
15 described above as preferred.

 The compounds of formula I-IV, including their salts, are also obtainable in the form of hydrates, or their crystals can include for example the solvent used for crystallization (present as solvates).

20 New starting materials and/or intermediates, as well as processes for the preparation thereof, are likewise the subject of this invention. In the preferred embodiment, such starting materials are used and reaction conditions so selected as to enable the preferred compounds to be
25 obtained.

 Starting materials of the invention, are known, are commercially available, or can be synthesized in analogy to or according to methods that are known in the art.


 In the preparation of starting materials, existing
30 functional groups which do not participate in the reaction should, if necessary, be protected. Preferred protecting groups, their introduction and their removal are described above or in the examples.

- 56 -

All remaining starting materials are known, capable of being prepared according to known processes, or commercially obtainable; in particular, they can be prepared using processes as described in the examples.

5 The following examples contain detailed descriptions of the methods of preparation of compounds of Formulas I-IV. These detailed descriptions fall within the scope, and serve to exemplify, the above described General Synthetic Procedures which form part of the invention. These detailed
10 descriptions are presented for illustrative purposes only and are not intended as a restriction on the scope of the invention.

 The compounds of this invention may contain one or more asymmetric centers and thus occur as racemates and
15 racemic mixtures, scalemic mixtures, single enantiomers, individual diastereomers and diastereomeric mixtures. All such isomeric forms of these compounds are expressly included in the present invention. The compounds of this invention may also be represented in multiple tautomeric
20 forms, for example, as illustrated below:



The invention expressly includes all tautomeric forms of the
25 compounds described herein. The compounds may also occur in cis- or trans- or E- or Z- double bond isomeric forms. All such isomeric forms of such compounds are expressly included in the present invention. All crystal forms of the compounds described herein are expressly included in the
30 present invention.

- 57 -

Substituents on ring moieties (e.g., phenyl, thienyl, etc.) may be attached to specific atoms, whereby they are intended to be fixed to that atom, or they may be drawn unattached to a specific atom, whereby they are intended to be attached at any available atom that is not already substituted by an atom other than H (hydrogen).

The compounds of this invention may contain heterocyclic ring systems attached to another ring system. Such heterocyclic ring systems may be attached through a carbon atom or a heteroatom in the ring system.

Alternatively, a compound of any of the formulas delineated herein may be synthesized according to any of the processes delineated herein. In the processes delineated herein, the steps may be performed in an alternate order and may be preceded, or followed, by additional protection/deprotection steps as necessary. The processes may further comprise use of appropriate reaction conditions, including inert solvents, additional reagents, such as bases (e.g., LDA, DIEA, pyridine, K_2CO_3 , and the like), catalysts, and salt forms of the above. The intermediates may be isolated or carried on *in situ*, with or without purification. Purification methods are known in the art and include, for example, crystallization, chromatography (liquid and gas phase), extraction, distillation, trituration, reverse phase HPLC and the like. Reactions conditions such as temperature, duration, pressure, and atmosphere (inert gas, ambient) are known in the art and may be adjusted as appropriate for the reaction.

As can be appreciated by the skilled artisan, the above synthetic schemes are not intended to comprise a comprehensive list of all means by which the compounds described and claimed in this application may be synthesized. Further methods will be evident to those of ordinary skill in the art. Additionally, the various

- 58 -

synthetic steps described above may be performed in an alternate sequence or order to give the desired compounds. Synthetic chemistry transformations and protecting group methodologies (protection and deprotection) useful in synthesizing the inhibitor compounds described herein are known in the art and include, for example, those such as described in R. Larock, *Comprehensive Organic Transformations*, VCH Publishers (1989); T.W. Greene and P.G.M. Wuts, *Protective Groups in Organic Synthesis*, 3rd. Ed., John Wiley and Sons (1999); L. Fieser and M. Fieser, *Fieser and Fieser's Reagents for Organic Synthesis*, John Wiley and Sons (1994); and L. Paquette, ed., *Encyclopedia of Reagents for Organic Synthesis*, John Wiley and Sons (1995).

The compounds of this invention may be modified by appending appropriate functionalities to enhance selective biological properties. Such modifications are known in the art and include those which increase biological penetration into a given biological compartment (e.g., blood, lymphatic system, central nervous system), increase oral availability, increase solubility to allow administration by injection, alter metabolism and alter rate of excretion.

Unless otherwise noted, all materials were obtained from commercial suppliers and used without further purification. All parts are by weight and temperatures are in Degrees centigrade unless otherwise indicated. All compounds showed NMR spectra consistent with their assigned structures.

In order that the invention described herein may be more readily understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

- 59 -

Analytical methods:

Unless otherwise indicated all HPLC analyses were run on an HP-1000 or HP-1050 system with an HP Zorbax SB-C₁₈ (5 μ) reverse phase column (4.6 x 150mm) run at 30 °C with a flow rate of 1.00 mL/min. The mobile phase used solvent A (H₂O/0.1% TFA) and solvent B (CH₃CN/0.1% TFA) with a 20 min gradient from 10% to 90% CH₃CN. The gradient was followed by a 2 min return to 10% CH₃CN and a 3 min flush. The peaks of interest eluted on the LC profiles at the times indicated.

LC-MS methods:**Method A:**

1. Samples were run on an HP-1100 system with an HP Zorbax SB-C₈ (5 μ) reverse phase column (4.6 x 50mm) run at 30°C with a flow rate of 0.75 mL/min.
2. The mobile phase used solvent A (H₂O/0.1% AcOH) and solvent B (CH₃CN/0.1% AcOH) with a 10 min gradient from 10% to 90% CH₃CN. The gradient was followed by a 1 min return to 10% CH₃CN and a 2 min flush.
3. The peaks of interest eluted on the LC profiles at the times indicated.

25

Method B:

4. Samples were run on an HP-1100 system with an HP Zorbax SB-C₈ (5 μ) reverse phase column (4.6 x 50mm) run at 30°C with a flow rate of 1.5 mL/min.
5. The mobile phase used solvent A (H₂O/0.1% AcOH) and solvent B (CH₃CN/0.1% AcOH) with a 5 min gradient from 10% to 90% CH₃CN. The gradient was followed by a 0.5 min return to 10% CH₃CN and a 1.5 min flush.

30

- 60 -

Preparative HPLC: Where indicated compounds of interest were purified via preparative HPLC using a Gilson workstation with a 30 x 100 mm column at 30 mL/min. The mobile phase used solvent A (H₂O/0.1% TFA) and solvent B (CH₃CN/0.1% TFA) with a 15 min gradient from 5% to 100% CH₃CN. The gradient was followed by a 2 min return to 5% CH₃CN.

Proton NMR Spectra:

Unless otherwise indicated all ¹H NMR spectra were run on an Varian series Mercury 300 or 400 MHz instrument. All observed protons are reported as parts per million (ppm) downfield from tetramethylsilane (TMS) or other internal reference in the appropriate solvent indicated.

15

The following abbreviations are used:


	AcOH, HOAc-	acetic acid
	CH ₃ CN -	acetonitrile
	ATP -	adenosine triphosphate
20	NH ₄ Cl	ammonium chloride
	NH ₄ OH -	ammonium hydroxide
	BINAP -	2,2'-bis(diphenylphosphino)-1,1'-binaphthyl
	BH ₃ -	borane
	BSA -	bovine serum albumin
25	DDQ -	2,3-dichloro-5,6-dicyano-1,4-benzoquinone
	CH ₂ Cl ₂ -	dichloromethane
	DIEA -	diisopropylethylamine
	DIAD -	diisopropyl azodicarboxylate
	EDC -	1-(3-dimethylaminopropyl)-3-
30		ethylcarbodiimide hydrochloride
	DMF -	dimethylformamide
	DMSO -	dimethyl sulfoxide
	DPPA -	diphenylphosphoryl azide

- 61 -

	DTT -	dithiothreitol
	EtOH -	ethanol
	EtOAc -	ethyl acetate
	Et ₂ O -	ethyl ether
5	FeSO ₄ -	ferric sulfate
	g -	gram
	h -	hour
	HBr -	hydrobromic acid
	HCl -	hydrochloric acid
10	H ₂ -	hydrogen
	HOBt -	hydroxybenzotriazole
	IPA -	isopropanol
	LAH -	lithium aluminum hydride
	LiOH -	lithium hydroxide
15	MgSO ₄ -	magnesium sulfate
	MnCl ₂ -	manganese chloride
	MeOH -	methanol
	MeI-	methyl iodide
	mg -	milligram
20	mL -	milliliter
	min -	minutes
	N ₂ -	nitrogen
	Pd/C-	palladium on carbon
	Pd(OAc) ₂ -	palladium acetate
25	Pd(PPh ₃) ₄ -	palladium tetrakis triphenylphosphine
	Pd ₂ (dba) ₃ -	tris(dibenzylideneacetone)di-palladium
	POCl ₃ -	phosphoryl chloride
	PCl ₅ -	phosphorous pentachloride
	P ₂ O ₅ -	phosphorous pentoxide
30	Pt/C -	platinum on carbon
	K ₂ CO ₃ -	potassium carbonate
	KOt-Bu -	potassium t-butoxide
	RT -	Room temperature
	NaHCO ₃ -	sodium bicarbonate

- 62 -

	Na ₂ CO ₃ -	sodium carbonate
	NaCl -	sodium chloride
	NaCN -	sodium cyanide
	NaCNBH ₃ -	sodium cyanoborohydride
5	NaH -	sodium hydride
	NaOH -	sodium hydroxide
	Na ₂ SO ₄ -	sodium sulfate
	NaOt-Bu -	sodium t-butoxide
	t-BuOH -	tert-butyl alcohol
10	t-BuOMe, MTBE -	tert-butylmethylether
	Boc -	tert-butyloxycarbonyl
	THF -	tetrahydrofuran
	TEA, Et ₃ N -	triethylamine
	TFA -	trifluoroacetic acid
15	PPh ₃ -	triphenyl phosphine
	H ₂ O -	water

Example 1

N⁴-(5-tert-Butylisoxazol-3-yl)-N²-(3,4,5-trimethoxy-phenyl)-pyrimidine-2,4-diamine

25 Step A

To a solution of 3-amino-5-*t*-butylisoxazole (922 mg 6.58 mmol) in 15 mL of DMF at 0 °C was added 268 mg (6.71 mmol) of NaH (60% dispersion in mineral oil) in 4 portions over 1 min. The resulting mixture was stirred at 0 °C for 30


- 63 -

min when 1.0 g (6.71 mmol) of 2,3-dichloropyrimidine was added in 1 portion. The resulting mixture was warmed to RT (ice melt in dewar) and was stirred at RT for 22 h. The reaction mixture was quenched with saturated aqueous NH_4Cl and diluted with H_2O and EtOAc . The organic layer was washed 1x with brine. The combined aqueous layer and brine wash were extracted 1x with EtOAc . The combined organics were dried over Na_2SO_4 , concentrated and purified by flash chromatography (elution with 2/1 hexanes/ EtOAc) to give (2-chloro-pyrimidin-4-yl)-(5-tert-butyl -isoxazol-3-yl)-amine as a yellowish white solid.

Step B

A mixture of (2-chloropyrimidin-4-yl)-(5-tert-butyl -isoxazol-3-yl)amine (50 mg 0.198 mmol Step A) and 3,4,5-trimethoxyaniline (36.2 mg 0.198 mmol) was suspended in 0.15 mL of a 2M solution of Et_3N -TFA in DMSO (stock solution prepared by combining 1.28 g of Et_3N -TFA and 3 mL of DMSO) in a sealed tube. The resulting mixture was heated at 100 °C for 4 h, cooled to RT, diluted with CH_2Cl_2 (attempted crystallization), concentrated and purified via preparative HPLC to give N^4 -(5-tert-butylisoxazol-3-yl)- N^2 -(3,4,5-trimethoxy-phenyl)-pyrimidine-2,4-diamine as a green-gray amorphous solid. MS m/z = 400. Calc'd for $\text{C}_{20}\text{H}_{25}\text{N}_5\text{O}_4$ - 399.19.

- 64 -

Example 2

5 ***N*⁴-(5-*t*-Butyl-isoxazol-3-yl)-*N*²-(4,5-dimethoxy-2-methyl-phenyl)-pyrimidine-2,4-diamine**

Step A: Preparation of (5-*t*-Butyl-isoxazol-3-yl)-(2-chloro-pyrimidin-4-yl)-amine

10 To a solution of 5-*t*-butyl-isoxazol-3-ylamine (10.0 g, 71.33 mmol) in *t*-BuOH (500 mL) at 20 °C was added NaOt-Bu (17.1 g, 178.3 mmol) in 3 portions over 10 min. The solution turned pale yellow with the appearance of a thick, white precipitate. The mixture was stirred at 20 °C for 1

15 h, then 2,4-dichloropyrimidine (15.9 g, 107.0 mmol) was added. The mixture was stirred for an additional 90 min. at 20 °C, accompanied by the dissolution of the white precipitate and appearance of an orange color. NH₄Cl (aq., sat., 100 mL) and H₂O (1 L) were added, and the mixture was

20 extracted with CH₂Cl₂ (3 x 200 mL). The combined organic layers were washed with H₂O (300 mL), brine, and dried with MgSO₄ before being filtered and concentrated under reduced pressure. Purification by chromatography of the crude reaction mixture (96:4 CH₂Cl₂:MeOH) provided the title

25 compound.

Step B: Preparation of *N*⁴-(5-*t*-Butyl-isoxazol-3-yl)-*N*²-(4,5-dimethoxy-2-methyl-phenyl)-pyrimidine-2,4-diamine

30 To a slurry of (5-*t*-butyl-isoxazol-3-yl)-(2-chloro-pyrimidin-4-yl)-amine (0.110 g, 0.435 mmol, Step A) in DMSO


- 65 -

(0.110 mL) was added 2-methyl-4,5-dimethoxyaniline (0.073 g, 0.435 mmol). The mixture was heated in a sealed tube at 110 °C for 90 min until TLC indicated the disappearance of starting materials. The mixture was taken up in CH₂Cl₂ (5 mL) followed by the addition of NaHCO₃ (aq., sat. 1 mL) and H₂O (5 mL). Extraction with CH₂Cl₂ (5 x 5 mL), followed by drying with MgSO₄ and filtration yielded a crude mixture that was concentrated under reduced pressure. Purification by chromatography (97:3 CH₂Cl₂:MeOH) on silica gel afforded the title compound as a purple solid. MS m/z = 384.2. Calc'd for C₂₀H₂₅N₅O₃: 383.45.

The following Examples were prepared from the corresponding amines in a manner similar to that described above for

Example 2:


Example 3



***N*⁴-(5-*t*-Butyl-isoxazol-3-yl)-*N*²-(1*H*-indazol-5-yl)-pyrimidine-2,4-diamine**

MS m/z = 350.3. Calc'd for C₁₈H₁₉N₇O: 349.40.


- 66 -

Example 4

5 ***N*⁴-(5-*t*-Butyl-isoxazol-3-yl)-*N*²-quinolin-3-yl-pyrimidine-
2,4-diamine**

MS *m/z* = 361.3. Calc'd for C₂₀H₂₀N₆O: 360.42.


10

Example 5

***N*⁴-(5-*t*-Butyl-isoxazol-3-yl)-*N*²-quinolin-6-yl-pyrimidine-
2,4-diamine**

15

MS *m/z* = 361.3. Calc'd for C₂₀H₂₀N₆O: 360.42.

Example 6

20


***N*⁴-(5-*t*-Butyl-isoxazol-3-yl)-*N*²-(1*H*-indazol-6-yl)-
pyrimidine-2,4-diamine**

- 67 -

MS m/z = 350.3. Calc'd for C₁₈H₁₉N₇O: 349.40.

Example 7

5



3-[4-(5-t-Butyl-isoxazol-3-ylamino)-pyrimidin-2-ylamino]- benzenesulfonamide

10

To a slurry of (5-t-butyl-isoxazol-3-yl)-(2-chloro-pyrimidin-4-yl)-amine (Example 2, Step A) (0.120 mg, 0.47 mmol) in DMSO (0.120 mL) was added 3-aminobenzenesulfonamide (0.082 g, 0.47 mmol). The reaction mixture was heated at 120 °C in a sealed tube for 1 h at which time TLC indicated the disappearance of starting material. IPA (5 mL) was added and the mixture was filtered and washed with IPA (2 x 1 mL) to yield the title compound as a white solid.





MS m/z = 389.3. Calc'd for C₁₇H₂₀N₆O₃S: 388.45.

20



Examples 8-13

Examples 8-13 were prepared from the corresponding amines in a manner similar to that described above for Example 7.

25


Example	Structure	Mass Calc'd	Mass Obs.
8	 <p>4-[4-(5-t-Butyl-isoxazol-3-ylamino)-pyrimidin-2-ylamino]-benzenesulfonamide</p>	C ₁₇ H ₂₀ N ₆ O ₃ S: 388.45	389.3
9	 <p>N⁴-(5-t-Butyl-isoxazol-3-yl)-N²-(3-chloro-phenyl)-pyrimidine-2,4-diamine</p>	C ₁₇ H ₁₈ ClN ₅ O: 343.82	344.2
10	 <p>N-{3-[4-(5-t-Butyl-isoxazol-3-ylamino)-pyrimidin-2-ylamino]-phenyl}-acetamide</p>	C ₁₉ H ₂₂ N ₆ O ₂ : 366.43	367.4
11	 <p>N-{4-[4-(5-t-Butyl-isoxazol-3-ylamino)-pyrimidin-2-ylamino]-phenyl}-N-methyl-acetamide</p>	C ₂₀ H ₂₄ N ₆ O ₂ : 380.45	381.4

- 69 -

12	 <p>N-Acetyl-4-[4-(5-<i>t</i>-butyl-isoxazol-3-ylamino)-pyrimidin-2-ylamino]-benzenesulfonamide</p>	$C_{19}H_{22}N_6O_4S:$ 430.49	431.4
13	 <p>N^4-(5-<i>t</i>-Butyl-isoxazol-3-yl)-N^2-(3-isopropyl-phenyl)-pyrimidine-2,4-diamine</p>	$C_{20}H_{25}N_5O:$ 351.46	352.3

- 70 -

Example 14



5 **N²-(3-Methoxy-4-morpholin-4-yl-phenyl)-N⁴-(5-phenyl-isoxazol-3-yl)-pyrimidine-2,4-diamine**

Step A: Preparation of (2-Chloro-pyrimidin-4-yl)-(5-phenyl-isoxazol-3-yl)-amine

10 The title compound was prepared in a manner similar to the method described in Example 2 (Step A) using the appropriate isoxazole reagent.

15 **Step B: Preparation of 4-(2-Methoxy-4-nitro-phenyl)-morpholine**

 To a mixture of 1-bromo-2-methoxy-4-nitro-benzene (2.5 g, 10.8 mmol), Pd₂(dba)₃ (0.124 g, 0.215 mmol), NaOt-Bu (1.55 g, 16.2 mmol) and BINAP (0.202 g, 0.323 mmol) in toluene (20 mL) at 20 °C was added morpholine (1.5 mL, 17.2 mmol) over 10 min. The mixture was stirred at 80 °C for 3 h when TLC indicated no starting material remained, at which time the mixture was evaporated under reduced pressure followed by the addition of H₂O (50 mL). CH₂Cl₂ extraction (3 x 15 mL), followed by drying of the combined organic layers with MgSO₄ afforded, after filtration and concentration, an orange oil. Chromatography on silica gel (97:3 CH₂Cl₂/MeOH) yielded pure compound.

- 71 -


Step C: Preparation of 3-Methoxy-4-morpholin-4-yl-phenylamine

To a solution of 4-(2-methoxy-4-nitro-phenyl)-morpholine (1.0 g, 4.2 mmol, Step B) in EtOH (25 mL) at 20 °C was added Pd/C (100 mg). The flask was capped with a rubber septum and H₂ pressure was applied through a balloon/needle. The reaction mixture was stirred at 20 °C for 12 h, filtered through sand/Celite® and concentrated. Purification by chromatography of the crude mixture (97:3 CH₂Cl₂:MeOH) afforded 3-methoxy-4-morpholin-4-yl-phenylamine as a purple solid.

Step D: Preparation of N²-(3-Methoxy-4-morpholin-4-yl-phenyl)-N⁴-(5-phenyl-isoxazol-3-yl)-pyrimidine-2,4-diamine

The title compound was prepared by the method described in Example 2, Step B using 3-methoxy-4-morpholin-4-yl-phenylamine (Step C). MS m/z = 445.3. Calc'd for C₂₄H₂₄N₆O₃: 444.50.

20

Example 15

25

N²-(3,4,5-Trimethoxyphenyl)-N⁴-(5-phenylisoxazol-3-yl)-2,4-pyrimidinediamine**Step A.**

n-Butyllithium (21.6 mL, 2.5M in hexane) was added dropwise to a solution of phenylacetylene (5 g, 0.05 mol) in

- 72 -

dry Et₂O (200 mL) maintaining at -78 °C under inert atmosphere. A solution of 2-pyridyl cyanate (7.2 g, 0.06 mol) in dry Et₂O (100 mL) was added dropwise keeping the temperature below -60 °C. The mixture was stirred for 30 min
5 at -78 °C then warmed to RT. The reaction was diluted with Et₂O and quenched with 1N NaOH. The organic layer was washed once with 1N HCl and once with saturated NaHCO₃. The solvent was removed under vacuum maintaining the temperature under 30 °C to give 3-phenyl-2-propynenitrile as a colorless
10 crystalline solid.

Step B.

To a stirred solution of 3-phenyl-2-propyne-nitrile (1.27 g, 0.01 mol, Step A) in EtOH (20 mL) was added at RT
15 hydroxylamine hydrochloride (3.5 g, 0.05 mol) followed by 10% NaOH (28 mL). The exothermic reaction was stirred at RT 24 h, then diluted with H₂O. The resulting precipitate was filtered off, washed with H₂O and dried over P₂O₅ to afford 3-amino-5-phenylisoxazole as a white crystalline solid.

20

Step C.


To a stirred suspension of 3-amino-5-phenylisoxazole (160 mg, 1 mmol, Step B) in *t*-BuOH (5 mL) was added NaO*t*-Bu (240 mg,). The mixture (slurry) was stirred at RT for 1 h.
25 Subsequently, 2,4-dichloro-1,3-pyrimidine (223 mg, 1.5 mmol) was added portion-wise and the reaction was stirred an additional 2 h. The reaction was quenched by addition of a saturated aqueous solution of NH₄Cl, and extracted with CH₂Cl₂. The extract was dried over MgSO₄ and evaporated. The
30 crude mixture was purified by flash chromatography using 20% EtOAc/Hexane as eluant to provide 2-chloro-N⁴-(5-phenylisoxazol-3-yl)-4-pyrimidineamine as a white solid.

- 73 -

Step D.

The title compound was prepared by the method described in Example 1 using 2-chloro-N⁴-(5-phenylisoxazol-3-yl)-4-pyrimidineamine and 3,4,5-trimethoxyaniline. MS

5 (MH⁺) = 420.4, Calc'd for C₂₂H₂₁N₅O₄ - 419.44.


Example 16

10

N²-(2-methyl-4,5-dimethoxyphenyl)-N⁴-(5-phenylisoxazol-3-yl)-2,4-pyrimidinediamine

The title compound was prepared by the method described in Example 1 using 2-chloro-N⁴-(5-phenylisoxazol-3-yl)-4-pyrimidineamine and 2-methyl-4,5-dimethoxyaniline. MS (MH⁺) = 404.7; Calc'd for C₂₂H₂₁N₅O₃ - 403.44.

15

Example 17

20

N²-(3,4,5-Trimethoxyphenyl)-N⁴-(3-cyclohexylisoxazol-5-yl)-2,4-pyrimidinediamine

- 74 -

Step A.

To 700 mL of THF cooled at -78 °C under inert atmosphere was added *n*-butyllithium (56 mL, 2.5 M in hexane) followed by CH₃CN (7.3 mL, 0.14 mol) maintaining the temperature below -70 °C. The mixture was stirred at -78 °C for 20 min then a solution of methyl cyclohexane carboxylate (10 g, 0.07 mol) in dry THF (300 mL) was added dropwise, maintaining the temperature under -60 °C. The reaction was stirred an additional 20 min at -78 °C, warmed to -20 °C and quenched by addition of saturated aq. NH₄Cl. The reaction mixture was diluted with H₂O and extracted with *t*-BuOMe. The solvents were removed under vacuum to give 3-cyclohexyl-3-oxopropanenitrile as a colorless oil.

Step B.

To a stirred solution of 3-cyclohexyl-3-oxopropanenitrile (1 g, 6.6 mmol, Step A) in EtOH (1 mL) at RT was added hydroxylamine (0.81 mL, 50% in H₂O). The reaction was stirred 12 h at RT. The solid was filtered off, washed with 1N NaOH, then H₂O and dried over P₂O₅ to give 5-amino-3-cyclohexylisoxazole.

Step C.

The 5-amino-3-cyclohexylisoxazole (Step B) was coupled to the 2,4-dichloropyridine using the general procedure described in Example 16 (step C) to afford 2-chloro-N⁴-(5-amino-3-cyclohexylisoxazolyl)-2,4-pyrimidineamine as an off white solid.

Step D.


The title compound was prepared by the method described in Example 1 using 2-chloro-N⁴-(3-cyclohexylisoxazol-5-yl)-4-pyrimidineamine (Step C) and

- 75 -

3,4,5-trimethoxyaniline. MS (MH^+) = 426.5; Calc'd for $C_{22}H_{27}N_5O_4$ - 425.49.

Example 18

5



N^2 -(4,5-Dimethoxy-2-methylphenyl)- N^4 -(5-adamantylisoxazol-3-yl)-2,4-pyrimidinediamine


10

The title compound was prepared by the method described in Example 1 using 2-chloro- N^4 -(5-adamantylisoxazol-3-yl)-4-pyrimidineamine and 2-methyl-4,5-dimethoxyaniline. MS (MH^+) = 462.2; Calc'd for $C_{26}H_{31}N_5O_3$ - 461.57.

15


Other compounds included in this invention are set forth in Tables 1-2 below.

Table 1.




#	R ²	R ¹
5	19. 1-benzylindazol-5-yl	5-tert-butyl-3-isoxazolyl
	20. 1-benzylindazol-5-yl	5-tert-butyl-3-isothiazolyl
	21. naphth-1-yl	5-tert-butyl-2-oxazolyl
	22. 4-benzylloxy-phenyl	5-tert-butyl-2-thiazolyl
	23. 4-benzylloxy-3-chlorophenyl	4-tert-butyl-2-thiazolyl
	24. 3-chloro-4-fluorophenyl	4-tert-butyl-2-oxazolyl
	25. 3-chloronaphth-1-yl	5-tert-butyl-1,3,4-oxadiazol-2-yl
	26. 4-methoxycarbonylphenyl	5-tert-butyl-1,3,4-thiadiazol-2-yl
	27. 3-(2-methoxyethoxy)-4-methoxyphenyl	5-tert-butyl-1,2,4-oxadiazol-2-yl
	28. 3,4-dimethoxyphenyl	5-tert-butyl-1,2,4-thiadiazol-2-yl
15	29. 3,4-diethoxyphenyl	2-tert-butyl-5-oxazolyl
	30. 3-bromophenyl	2-tert-butyl-5-thiazolyl
	31. 3-methoxy-4-(2-methoxyethoxy)phenyl	2-tert-butyl-4-thiazolyl
	32. 3-methylphenyl	2-tert-butyl-4-oxazolyl
	33. 4,5-dimethoxy-3-(2-morpholin-4-yl-ethoxy)phenyl	5-tert-butyl-1,3,4-oxadiazol-2-yl
	34. 3-aminocarbonylphenyl	5-tert-butyl-1,2,4-oxadiazol-2-yl

Table 1. cont.




	<u>R</u>	
	<u>R²</u>	<u>R³</u>
5	35. 3-CH ₃ OCOCH ₂ -phenyl	5-tert-butyl-1,3,4-thiadiazol-2-yl
	36. 3-(4-nitrophenylsulfonylamino)phenyl	5-tert-butyl-1,3,4-thiadiazol-2-yl
	37. 5-chloro-2-methyl-phenyl	5-tert-butyl-3-isoxazolyl
	38. 4-[HO(CH ₂) ₂ O]phenyl	5-tert-butyl-3-isothiazolyl
	39. 6-indazolyl	2-tert-butyl-4-thiazolyl
10	40. 5-benzimidazolyl	2-tert-butyl-4-oxazolyl
	41. 3,4-dimethoxy-6-methylphenyl	5-tert-butyl-3-isothiazolyl

Table 2.




#	R ²	R ³
42.	3-quinolyl	tert-butyl
43.	6-quinolyl	tert-butyl
44.	2,5-dimethoxyphenyl	tert-butyl
45.	2-chloro-5-methoxyphenyl	tert-butyl
46.	4-methoxy-2-methylphenyl	tert-butyl
47.	5-methoxy-2-methylphenyl	tert-butyl
48.	3-trifluoromethoxyphenyl	tert-butyl
49.	3-(1,1,2,2-tetrafluoroethoxy)phenyl	tert-butyl
50.	4-trifluoromethoxyphenyl	tert-butyl
51.	6-indazolyl	tert-butyl
52.	5-benzimidazolyl	tert-butyl
53.	1-methyl-5-indazolyl	isopropyl
54.	2-methyl-2H-indazol-5-yl	tert-butyl
55.	3-isopropylphenyl	tert-butyl
56.	3,5-dimethoxyphenyl	tert-butyl
57.	3,4-dimethoxyphenyl	tert-butyl
58.	1-methoxy-3-naphthyl	tert-butyl

Table 2. cont.




5	#	R ²	R ³
	59.	1,3-benzodioxol-5-yl	tert-butyl
	60.	2-methoxyphenyl	isopropyl
	61.	2-methoxyphenyl	tert-butyl
	62.	3-methoxyphenyl	tert-butyl
10	63.	4-methoxyphenyl	tert-butyl
	64.	3,4-diethoxyphenyl	tert-butyl
	65.	3,5-bis(trifluoromethyl)phenyl	tert-butyl
	66.	3-fluoro-5-(trifluoromethyl)phenyl	tert-butyl
	67.	3-ethylphenyl	tert-butyl
15	68.	4-ethylphenyl	tert-butyl
	69.	3-ethynylphenyl	tert-butyl
	70.	3-cyanophenyl	tert-butyl
	71.	2-cyano-4,5-dimethoxyphenyl	tert-butyl
	72.	4-cyanophenyl	tert-butyl
20	73.	4-aminocarbonyl-3-methoxyphenyl	tert-butyl
	74.	4-(N-hydroxyethylamino)carbonyl-3-methoxyphenyl	tert-butyl
	75.	3-methoxy-4-pentafluoroethylphenyl	tert-butyl
	76.	2-methoxycarbonyl-4,5-dimethoxyphenyl	tert-butyl
	77.	6-methoxy-8-quinolyl	tert-butyl

Table 2. cont.




#	R ²	R ³
5		
78.	2-(3,4,5-trimethoxyphenyl)	adamantyl
79.	4-(3-Dimethylamino-propoxy)-phenyl	tert-butyl
80.	3,5-Dimethoxy-4-[2-(4-methyl-piperazin-1-yl)ethoxy]phenyl	phenyl
81.	3,5-Dimethoxy-4-[3-(4-methyl-piperazin-1-yl)propoxy]phenyl	cyclohexyl
82.	3-Methoxy-4-[3-(4-methyl-piperazin-1-yl)-propoxy]phenyl	tert-butyl
83.	3-Methoxy-4-[2-(4-methyl-piperazin-1-yl)-ethoxy]phenyl	adamantyl
84.	2-Methyl-4-(4-methyl-piperazin-1-yl)-phenyl	tert-butyl
85.	4-(4-Isopropyl-piperazin-1-yl)-phenyl	cyclohexyl
86.	3-Fluoro-4-(4-methyl-piperazin-1-yl)-phenyl	phenyl
87.	3-Fluoro-4-(3-piperidin-1-yl-propoxy)-phenyl	adamantyl
88.	3,4-Dimethoxy-5-[3-(4-methyl-piperazin-1-yl)-propoxy]-phenyl	cyclohexyl
89.	3-Methoxy-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl	tert-butyl
90.	3-Methoxy-4-[2-(1-methyl-piperidin-4-yl)-ethoxy]-phenyl	phenyl
91.	4-(1-tert-butoxycarbonyl-piperazin-4-yl)phenyl	adamantyl
92.	4-(4-piperazinyl)phenyl	cyclohexyl
93.	4-(1-tert-butoxycarbonyl-piperazin-4-yl)- 3-difluoromethoxy-phenyl	tert-butyl
10		
15		
20		

Table 2. cont.




#	R ²	R ³
94.	3-Fluoro-4-(2-piperidin-1-yl-ethoxy)-phenyl	phenyl
95.	3-Fluoro-4-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl	adamantyl
96.	3-Fluoro-4-[3-(4-methyl-piperazin-1-yl)-propoxy]-phenyl	cyclohexyl
97.	4-[2-(4-Methyl-piperazin-1-yl)-ethoxy]-phenyl	tert-butyl
98.	3,5-Dimethoxy-4-(2-morpholin-4-yl-ethoxy)-phenyl	phenyl
99.	3,5-Dimethoxy-4-(2-piperidin-1-yl-ethoxy)-phenyl	adamantyl
100.	2-Methyl-4-(3-piperidin-1-yl-propoxy)-phenyl	cyclohexyl
101.	2-Methyl-4-[3-(4-methyl-piperazin-1-yl)-propoxy]-phenyl	tert-butyl
102.	2-Fluoro-4,5-dimethoxy-phenyl	phenyl
103.	2-anthracenyl	tert-butyl
104.	2-benzimidazolyl	phenyl
105.	Phenanthren-3-yl	cyclohexyl
106.	7-HOSO ₂ -naphth-2-yl	adamantyl
107.	3-carbazolyl	tert-butyl
108.	2-phenanthrenyl	phenyl
109.	5-HOSO ₂ -2-naphthyl	cyclohexyl
110.	8-HOSO ₂ -2-naphthyl	adamantyl

Table 2. cont.




5	#	R ²	R ³
	111.	2-HOSO ₂ -6-Naphthyl	tert-butyl
	112.	6-(4-Chlorophenoxy) Pyridin-3-yl	phenyl
	113.	6-(3-Pyridyloxy) Pyridin-3-yl	adamantyl
	114.	6-(4-Chloro-2-Cyclohexylphenoxy) Pyridin-3-yl	cyclohexyl
10	115.	2-(2-Pyridyl) Benzimidazol-5-yl	tert-butyl
	116.	2-dibenzofuryl	adamantyl
	117.	2-Phenylbenzoxazol-6-yl	phenyl
	118.	2,3-bis(4-methoxyphenyl) quinoxalin-6-yl	cyclohexyl
	119.	2-dibenzofuryl	tert-butyl
15	120.	1,1-dioxobenzo[b]thiophen-6-yl	cyclohexyl
	121.	1-[(4-Methylphenyl) sulfonyl]-1H-indol-5-yl	phenyl

Table 3.



5	#	R ²	R ³
10	122.	3-quinolyl	tert-butyl
	123.	6-quinolyl	tert-butyl
	124.	2,5-dimethoxyphenyl	tert-butyl
	125.	2-chloro-5-methoxyphenyl	tert-butyl
	126.	4-methoxy-2-methylphenyl	phenyl
15	127.	5-methoxy-2-methylphenyl	cyclohexyl
	128.	3-trifluoromethoxyphenyl	adamantyl
	129.	3-(1,1,2,2-tetrafluoroethoxy)phenyl	tert-butyl
	130.	4-trifluoromethoxyphenyl	phenyl
	131.	6-indazolyl	cyclohexyl
20	132.	5-benzimidazolyl	tert-butyl
	133.	1-methyl-5-indazolyl	isopropyl
	134.	2-methyl-2H-indazol-5-yl	phenyl
	135.	3-isopropylphenyl	cyclohexyl
	136.	3,5-dimethoxyphenyl	adamantyl
	137.	3,4-dimethoxyphenyl	tert-butyl
	138.	1-methoxy-3-naphthyl	phenyl
	139.	1-(4-Methylpiperazino)isoquinolin-3-yl	cyclohexyl

Table 3.



#	R ²	R ³
140.	2-Phenyl-6-quinoxaliny1	adamantyl
141.	2-(4-Dimethylamino-phenyl)-benzooxazol-5-yl	tert-butyl
142.	2-Phenyl-3H-benzimidazol-5-yl	phenyl
143.	1-(2-Hydroxyethyl)-2-methyl-1H-benzimidazol-5-yl	cyclohexyl
144.	[4,5-dihydro-4-carboxy-thiazol-2-yl]-benzothiazol-6-yl	adamantyl
145.	3-(Aminocarbonyl)-1H-indol-5-yl	tert-butyl

- 85 -

Although the pharmacological properties of the compounds of Formulas I-IV vary with structural change, in general, activity possessed by compounds of Formulas I-IV may be demonstrated in vivo. The pharmacological properties of the compounds of this invention may be confirmed by a number of pharmacological in vitro assays. The exemplified pharmacological assays which follow have been carried out with the compounds according to the invention and their derivatives.

10

BIOLOGICAL EVALUATION

Kinase Inhibition

15 The compounds described herein are screened in the following manner. Kinases suitable for use in the following protocol to determine kinase activity of the compounds described herein include, but are not limited to: Lck, Lyn, Src, Fyn, Syk, Zap-70, Itk, Tec, Btk, EGFR, ErbB2, Kdr, Flt-
20 1, Flt-3, Tek, c-Met, InsR, and AKT.

Kinases are expressed as either kinase domains or full length constructs fused to glutathione S-transferase (GST) or polyHistidine tagged fusion proteins in either E. coli or Baculovirus-High Five expression systems. They are purified
25 to near homogeneity by affinity chromatography essentially as previously described (Lehr et al., 1996; Gish et al., 1995). In some instances, kinases are co-expressed or mixed with purified or partially purified regulatory polypeptides prior to measurement of activity.

30 Kinase activity and inhibition are measured essentially by established protocols (Braunwalder et al., 1996). Briefly, the transfer of $^{33}\text{PO}_4$ from ATP to the synthetic substrates poly(Glu, Tyr) 4:1 or poly(Arg, Ser) 3:1 attached to the bioactive surface of microtiter plates

- 86 -

serves as the basis to evaluate enzyme activity. After an incubation period, the amount of phosphate transferred is measured by first washing the plate with 0.5% phosphoric acid, adding liquid scintillant, and then counting in a liquid scintillation detector. The IC_{50} is determined by the concentration of compound that causes a 50% reduction in the amount of ^{33}P incorporated onto the substrate bound to the plate.

Other similar methods whereby phosphate is transferred to peptide or polypeptide substrate containing tyrosine, serine, threonine, or histidine, either alone, in combination, or in combination with other amino acids, in solution or immobilized (i.e., solid phase) are also useful. For example, transfer of phosphate to a peptide or polypeptide can also be detected using scintillation proximity (Wu et al., 2000), ELISA (Cleaveland et al., 1990), Fluorescence Polarization (Seethala and Menzel, 1998), and homogeneous time-resolved fluorescence (HTRF, Kolb et al., 1998). Alternatively, kinase activity can be measured using antibody-based methods whereby an antibody or polypeptide is used as a reagent to detect phosphorylated target polypeptide. Compounds of the present invention showed inhibition of IGF-1R kinase at doses less than 50 μM .

The compounds of examples 1-3, and 14-18 inhibited IGF-1R kinase at a level below 100 nM (IC_{50}).

References:

- Braunwalder et al. (1996). *Anal. Biochem.* 234(1):23-26.
- Cleaveland et al. (1990). *Anal Biochem.* 190(2):249-53.
- Gish et al. (1995). *Protein Eng.* 8(6):609-614.
- Kolb et al. (1998). *Drug Discov. Today.* 3:333-342.
- Lehr et al. (1996). *Gene* 169(2):27527-9.

- 87 -

Seethala et al. (1998). *Anal Biochem.* 255(2):257-62.

Wu et al. (2000). *Comb Chem High Throughput Screen.*
3(1):27-36.

5

IGF-1R assay summary protocols

IGF-1-induced DNA synthesis.

- 10 Human tumor cell lines or a rat fibroblast cell line are
plated out in flat-well plates in complete medium and
allowed to adhere overnight. The cells are then starved in
medium containing 0.1% bovine serum albumin (BSA) overnight,
pre-incubated for 1 h with or without dilutions of compound,
15 then activated overnight with 50 ng/mL insulin-like growth
factor (IGF-1). Proliferation is determined by the level of
³H-thymidine incorporation into DNA. IC₅₀'s are determined
by comparing the level of thymidine incorporation found in
the presence of compound compared to controls.
20 Examples 14-15 and 17-18 inhibited 3T3 proliferation at a
level below 150 nM.

IGF-1R auto-phosphorylation.

- 25 Murine fibroblast cells stably transfected with the
human IGF-1R are plated out in flat-well plates in complete
media and allowed to adhere overnight. The cells are then
starved in medium containing 0.1% bovine serum albumin, pre-
incubated with or without dilutions of compound, then
30 activated for 5 min with 100 ng/mL IGF-1. The cells are
lysed and proteins are separated by SDS-PAGE. The level of
phosphotyrosine on IGF-1R β -chain is determined by western
blotting with an anti-phospho-IGF-1R β -specific antibody.
IC₅₀'s are determined by comparing the level of
35 phosphotyrosine found in the presence of compound compared
to controls.

- 88 -

Representative compounds tested under the following example protocols exhibit cellular activities consistent with their observed enzyme inhibition activities.

5 Formulations

Also embraced within this invention is a class of pharmaceutical compositions comprising the active compounds of Formula I in association with one or more non-toxic, pharmaceutically-acceptable carriers and/or diluents and/or adjuvants (collectively referred to herein as "carrier" materials) and, if desired, other active ingredients. The active compounds of the present invention may be administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended. The compounds and compositions of the present invention may, for example, be administered orally, mucosally, topically, rectally, pulmonarily such as by inhalation spray, or parentally including intravascularly, intravenously, intraperitoneally, subcutaneously, intramuscularly intrasternally and infusion techniques, in dosage unit formulations containing conventional pharmaceutically acceptable carriers, adjuvants, and vehicles.

25 The pharmaceutically active compounds of this invention can be processed in accordance with conventional methods of pharmacy to produce medicinal agents for administration to patients, including humans and other mammals.

30 For oral administration, the pharmaceutical composition may be in the form of, for example, a tablet, capsule, suspension or liquid. The pharmaceutical composition is preferably made in the form of a dosage unit containing a particular amount of the active ingredient.

35 Examples of such dosage units were tablets or capsules. For

- 89 -

example, these may contain an amount of active ingredient from about 1 to 2000 mg, preferably from about 1 to 500 mg, more preferably from about 5 to 150 mg. A suitable daily dose for a human or other mammal may vary widely depending
5 on the condition of the patient and other factors, but, once again, can be determined using routine methods.

The amount of compounds which were administered and the dosage regimen for treating a disease condition with the compounds and/or compositions of this invention depends on a
10 variety of factors, including the age, weight, sex and medical condition of the subject, the type of disease, the severity of the disease, the route and frequency of administration, and the particular compound employed. Thus, the dosage regimen may vary widely, but can be determined
15 routinely using standard methods. A daily dose of about 0.01 to 500 mg/kg body weight, preferably between about 0.5 and about 50 mg/kg body weight and most preferably between about 0.1 to 20 mg/kg body weight, may be appropriate may be appropriate. The daily dose can be administered in one to
20 four doses per day.

For therapeutic purposes, the active compounds of this invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered per os, the compounds may be admixed with
25 lactose, sucrose, starch powder, cellulose esters of alkanolic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl
30 alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets may contain a controlled-release formulation as may be provided in a dispersion of active compound in hydroxypropylmethyl cellulose.

- 90 -

In the case of psoriasis and other skin conditions, it may be preferable to apply a topical preparation of compounds of this invention to the affected area two to four times a day.

5 Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin (e.g., liniments, lotions, ointments, creams, or pastes) and drops suitable for administration to the eye, ear, or nose. A suitable topical
10 dose of active ingredient of a compound of the invention is 0.1 mg to 150 mg administered one to four, preferably one or two times daily. For topical administration, the active ingredient may comprise from 0.001% to 10% w/w, e.g., from 1% to 2% by weight of the formulation, although it may
15 comprise as much as 10% w/w, but preferably not more than 5% w/w, and more preferably from 0.1% to 1% of the formulation.

When formulated in an ointment, the active ingredients may be employed with either paraffinic or a water-miscible ointment base. Alternatively, the active ingredients may be
20 formulated in a cream with an oil-in-water cream base. If desired, the aqueous phase of the cream base may include, for example at Least 30% w/w of a polyhydric alcohol such as propylene glycol, butane-1,3-diol, mannitol, sorbitol, glycerol, polyethylene glycol and mixtures thereof. The
25 topical formulation may desirably include a compound which enhances absorption or penetration of the active ingredient through the skin or other affected areas. Examples of such dermal penetration enhancers include dimethylsulfoxide and related analogs.

30 The compounds of this invention can also be administered by a transdermal device. Preferably transdermal administration will be accomplished using a patch either of the reservoir and porous membrane type or of a solid matrix variety. In either case, the active agent is delivered

- 91 -

continuously from the reservoir or microcapsules through a membrane into the active agent permeable adhesive, which is in contact with the skin or mucosa of the recipient. If the active agent is absorbed through the skin, a controlled and
5 predetermined flow of the active agent is administered to the recipient. In the case of microcapsules, the encapsulating agent may also function as the membrane.

The oily phase of the emulsions of this invention may be constituted from known ingredients in a known manner.
10 While the phase may comprise merely an emulsifier, it may comprise a mixture of at least one emulsifier with a fat or an oil or with both a fat and an oil. Preferably, a hydrophilic emulsifier is included together with a lipophilic emulsifier which acts as a stabilizer. It is also
15 preferred to include both an oil and a fat. Together, the emulsifier(s) with or without stabilizer(s) make-up the so-called emulsifying wax, and the wax together with the oil and fat make up the so-called emulsifying ointment base which forms the oily dispersed phase of the cream
20 formulations. Emulsifiers and emulsion stabilizers suitable for use in the formulation of the present invention include Tween 60, Span 80, cetostearyl alcohol, myristyl alcohol, glyceryl monostearate, sodium lauryl sulfate, glyceryl distearate alone or with a wax, or other materials well
25 known in the art.

The choice of suitable oils or fats for the formulation is based on achieving the desired cosmetic properties, since the solubility of the active compound in most oils likely to be used in pharmaceutical emulsion
30 formulations is very low. Thus, the cream should preferably be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or other containers. Straight or branched chain, mono- or dibasic alkyl esters such as di-isoadipate, isocetyl stearate,

- 92 -

propylene glycol diester of coconut fatty acids, isopropyl myristate, decyl oleate, isopropyl palmitate, butyl stearate, 2-ethylhexyl palmitate or a blend of branched chain esters may be used. These may be used alone or in
5 combination depending on the properties required. Alternatively, high melting point lipids such as white soft paraffin and/or liquid paraffin or other mineral oils can be used.

Formulations suitable for topical administration to
10 the eye also include eye drops wherein the active ingredients were dissolved or suspended in suitable carrier, especially an aqueous solvent for the active ingredients. The active ingredients were preferably present in such formulations in a concentration of 0.5 to 20%,
15 advantageously 0.5 to 10% and particularly about 1.5% w/w.

Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules
20 using one or more of the carriers or diluents mentioned for use in the formulations for oral administration or by using other suitable dispersing or wetting agents and suspending agents. The compounds may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil,
25 cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, tragacanth gum, and/or various buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art. The active ingredient may also be administered by injection as a
30 composition with suitable carriers including saline, dextrose, or water, or with cyclodextrin (i.e. Captisol), cosolvent solubilization (i.e. propylene glycol) or micellar solubilization (i.e. Tween 80).

- 93 -

The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles
5 and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland
10 fixed oil may be employed, including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

For pulmonary administration, the pharmaceutical composition may be administered in the form of an aerosol or with an inhaler including dry powder aerosol.

15 Suppositories for rectal administration of the drug can be prepared by mixing the drug with a suitable non-irritating excipient such as cocoa butter and polyethylene glycols that are solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the
20 rectum and release the drug.

The pharmaceutical compositions may be subjected to conventional pharmaceutical operations such as sterilization and/or may contain conventional adjuvants, such as preservatives, stabilizers, wetting agents, emulsifiers,
25 buffers etc. Tablets and pills can additionally be prepared with enteric coatings. Such compositions may also comprise adjuvants, such as wetting, sweetening, flavoring, and perfuming agents.

Pharmaceutical compositions of this invention comprise
30 a compound of the formulas described herein or a pharmaceutically acceptable salt thereof; an additional agent selected from a kinase inhibitory agent (small molecule, polypeptide, antibody, etc.), an immunosuppressant, an anticancer agent, an anti-viral agent,

- 94 -

antiinflammatory agent, antifungal agent, antibiotic, or an anti-vascular hyperproliferation compound; and any pharmaceutically acceptable carrier, adjuvant or vehicle. Alternate compositions of this invention comprise a compound
5 of the formulae described herein or a pharmaceutically acceptable salt thereof; and a pharmaceutically acceptable carrier, adjuvant or vehicle. Such compositions may optionally comprise one or more additional therapeutic agents, including, for example, kinase inhibitory agents
10 (small molecule, polypeptide, antibody, etc.), immunosuppressants, anti-cancer agents, anti-viral agents, antiinflammatory agents, antifungal agents, antibiotics, or anti-vascular hyperproliferation compounds.

The term "pharmaceutically acceptable carrier or
15 adjuvant" refers to a carrier or adjuvant that may be administered to a patient, together with a compound of this invention, and which does not destroy the pharmacological activity thereof and is nontoxic when administered in doses sufficient to deliver a therapeutic amount of the compound.

20 Pharmaceutically acceptable carriers, adjuvants and vehicles that may be used in the pharmaceutical compositions of this invention include, but were not limited to, ion exchangers, alumina, aluminum stearate, lecithin, self-emulsifying drug delivery systems (SEDDS) such as d-a-
25 tocopherol polyethyleneglycol 1000 succinate, surfactants used in pharmaceutical dosage forms such as Tweens or other similar polymeric delivery matrices, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride
30 mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances,

- 95 -

polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat. Cyclodextrins such as α -, β -, and γ -cyclodextrin, or chemically modified
5 derivatives such as hydroxyalkylcyclodextrins, including 2- and 3-hydroxypropyl-cyclodextrins, or other solubilized derivatives may also be advantageously used to enhance delivery of compounds of the formulae described herein.

The pharmaceutical compositions of this invention may
10 be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, emulsions and aqueous suspensions, dispersions and solutions. In the case of tablets for oral use, carriers which were commonly used include lactose and corn starch. Lubricating agents,
15 such as magnesium stearate, were also typically added. For oral administration in a capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions and/or emulsions were administered orally, the active ingredient may be suspended or dissolved in an oily
20 phase is combined with emulsifying and/or suspending agents. If desired, certain sweetening and/or flavoring and/or coloring agents may be added.

The pharmaceutical compositions of this invention may comprise formulations utilizing liposome or
25 microencapsulation techniques. Such techniques were known in the art.

The pharmaceutical compositions of this invention may be administered by nasal aerosol or inhalation. Such compositions were prepared according to techniques well-
30 known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art.

- 96 -

The specification and claims contain listing of species using the language "selected from . . . and . . ." and "is . . . or . . ." (sometimes referred to as Markush groups). When this language is used in this application, 5 unless otherwise stated it is meant to include the group as a whole, or any single members thereof, or any subgroups thereof. The use of this language is merely for shorthand purposes and is not meant in any way to limit the removal of individual elements or subgroups as needed.

10 The foregoing is merely illustrative of the invention and is not intended to limit the invention to the disclosed compounds. Variations and changes which are obvious to one skilled in the art are intended to be within the scope and nature of the invention which are defined in the appended 15 claims.

From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the 20 invention to adapt it to various usages and conditions.


All mentioned references, patents, applications and publications, are hereby incorporated by reference in their entirety, as if here written.

- 97 -

What is Claimed is:

1. A compound of Formula I

5



wherein R¹ is heteroaryl containing at least one oxygen or sulfur heteroatom, wherein R¹ is optionally substituted with 1-4 substituents independently selected from R³;

10 wherein R² is selected from

H,

C₁₋₁₀-alkyl,

C₂₋₁₀-alkenyl,

C₂₋₁₀-alkynyl,

15

C(O)R⁵,

COOR⁵,

C(O)NR⁵R⁵,

S(O)_nR⁵,

C₃₋₁₀-cycloalkyl,

20

C₄₋₁₀-cycloalkenyl,

aryl optionally substituted with 1-5 substituents independently selected from R³,

R⁴,

C₁₋₁₀-alkyl substituted with 1-3 substituents

25

independently selected from aryl, R⁷ and R⁴,

C₃₋₁₀-cycloalkyl substituted with 1-3 substituents

independently selected from aryl, R⁷ and R⁴, and

C₂₋₁₀-alkenyl substituted with 1-3 substituents

independently selected from aryl, R⁷ and R⁴;

30 wherein R³ is independently selected from

H,

C₁₋₁₀-alkyl,

- 98 -

- C₂₋₁₀-alkenyl,
 C₂₋₁₀-alkynyl,
 C₃₋₁₀-cycloalkyl,
 C₄₋₁₀-cycloalkenyl,
 5 aryl,
 R⁴,
 halo,
 SR⁵,
 OR⁵,
 10 OC(O)R⁵,
 NR⁵R⁵,
 NR⁵R⁶,
 COOR⁵,
 nitro,
 15 cyano,
 C(O)R⁵,
 C(O)C(O)R⁵,
 C(O)NR⁵R⁵,
 S(O)_nR⁵,
 20 S(O)_nNR⁵R⁵,
 NR⁵C(O)NR⁵R⁵,
 NR⁵C(O)C(O)R⁵,
 NR⁵C(O)R⁵,
 NR⁵(COOR⁵),
 25 NR⁵C(O)R⁴,
 NR⁵S(O)_nNR⁵R⁵,
 NR⁵S(O)_nR⁵,
 NR⁵S(O)_nR⁴,
 NR⁵C(O)C(O)NR⁵R⁵,
 30 NR⁵C(O)C(O)NR⁵R⁶,
 C₁₋₁₀-alkyl substituted with 1-3 substituents
 independently selected from aryl, R⁷ and R⁴; and
 C₂₋₁₀-alkenyl substituted with 1-3 substituents
 independently selected from aryl, R⁷ and R⁴;

- 99 -

wherein R^4 is independently a 5-8 membered monocyclic, 8-12 membered bicyclic, or 11-14 membered tricyclic saturated, partially saturated or unsaturated ring system comprising 1-3 heteroatoms if monocyclic, 1-6 heteroatoms if bicyclic, or 1-9 heteroatoms if tricyclic, said heteroatoms independently selected from O, N, or S, which may be saturated or unsaturated, and wherein 0, 1, 2 or 3 atoms of each ring may be substituted by a substituent independently selected from C_{1-10} -alkyl, C_{2-10} -alkenyl, C_{2-10} -alkynyl, C_{3-10} -cycloalkyl, C_{4-10} -cycloalkenyl, halo, haloalkyl, sulfo, oxo, SR^5 , OR^5 , NR^5R^5 , NR^5R^6 , NR^6R^6 , $COOR^5$, nitro, cyano, $S(O)_nR^5$, $S(O)_nNR^5R^5$, $C(O)R^5$ and $C(O)NR^5R^5$;

wherein R^5 is independently selected from H, C_{1-10} -alkyl, C_{2-10} -alkenyl, C_{2-10} -alkynyl, C_{3-10} -cycloalkyl, C_{4-10} -cycloalkenyl, R^4 , aryl optionally substituted with 1-3 substituents independently selected from R^3 , $C_{1-C_{10}}$ alkyl substituted with 1-3 substituents independently selected from aryl, R^7 and R^4 ; $C_{3-C_{10}}$ cycloalkyl substituted with 1-3 substituents independently selected from aryl, R^7 and R^4 ; and $C_{2-C_{10}}$ alkenyl substituted with 1-3 substituents independently selected from aryl, R^7 and R^4 ;

wherein R^6 is selected from $-C(O)R^5$, $-COOR^5$, $-C(O)NR^5R^5$ and $-S(O)_nR^5$;

wherein R^7 is independently halo, $-CF_3$, $-SR^5$, $-OR^5$, $-OC(O)R^5$, $-NR^5R^5$, $-NR^5R^6$, $-NR^6R^6$, $-COOR^5$, $-NO_2$, $-CN$, $-C(O)R^5$, $OC(O)NR^5R^5$, $-C(O)NR^5R^5$, $-N(R^5)C(O)R^5$, $-N(R^5)(COOR^5)$ and $-S(O)_nNR^5R^5$; and

wherein n is 1 or 2;

and pharmaceutically acceptable derivative thereof.

2. The compound of claim 1, wherein R^1 is 5- or 6-membered heteroaryl containing at least one oxygen or

- 100 -

sulfur heteroatom, wherein R^1 is optionally substituted
with 1-4 substituents independently selected from R^3 ;
wherein R^2 is selected from

- C₁₋₆-alkyl,
- 5 C₂₋₆-alkenyl,
- C₂₋₆-alkynyl,
- C₃₋₆-cycloalkyl,
- C₄₋₆-cycloalkenyl,
- R⁴,
- 10 phenyl optionally substituted with 1-4 substituents
independently selected from R^3 ,
- C₁₋₆-alkyl substituted with 1-3 substituents independently
selected from aryl, R^7 and R^4 ,
- C₃₋₆-cycloalkyl substituted with 1-3 substituents
- 15 independently selected from aryl, R^7 and R^4 , and
- C₂₋₆-alkenyl substituted with 1-3 substituents
independently selected from aryl and R^4 ,
- wherein R^3 is independently selected from
- C₁₋₆-alkyl,
- 20 C₂₋₆-alkenyl,
- C₂₋₆-alkynyl,
- C₃₋₆-cycloalkyl,
- C₄₋₆-cycloalkenyl,
- phenyl,
- 25 R⁴,
- halo,
- SR⁵,
- OR⁵,
- OC(O)R⁵,
- 30 NR⁵R⁵,
- NR⁵R⁶,
- COOR⁵,
- NO₂,
- CN,

- 101 -

- $C(O)R^5$,
 $C(O)C(O)R^5$,
 $C(O)NR^5R^5$,
 $S(O)_nR^5$,
5 $S(O)_nNR^5R^5$,
 $NR^5C(O)NR^5R^5$,
 $NR^5C(O)C(O)R^5$,
 $NR^5C(O)R^5$,
 $NR^5(COOR^5)$,
10 $NR^5C(O)R^4$,
 $NR^5S(O)_nNR^5R^5$,
 $NR^5S(O)_nR^5$,
 $NR^5S(O)_nR^4$,
 $NR^5C(O)C(O)NR^5R^5$,
15 $NR^5C(O)C(O)NR^5R^6$,
 C_{1-6} -alkyl substituted with 1-3 substituents independently
selected from aryl, R^7 and R^4 ; and
 C_{2-6} -alkenyl substituted with 1-3 substituents
independently selected from aryl, R^7 and R^4 ;
20 wherein R^4 is independently a 5-7 membered monocyclic, or 8-
11 membered bicyclic saturated, partially saturated or
unsaturated ring system comprising 1-3 heteroatoms if
monocyclic, or 1-6 heteroatoms if bicyclic, said
heteroatoms independently selected from O, N, or S, which
25 may be saturated or unsaturated, and wherein 0, 1, 2 or 3
atoms of each ring may be substituted by a substituent
independently selected from C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -
alkynyl, halo, C_{1-6} -haloalkyl, oxo, SR^5 , OR^5 , NR^5R^5 , $COOR^5$,
nitro, cyano, $S(O)_nR^5$, $S(O)_nNR^5R^5$, $C(O)R^5$ and $C(O)NR^5R^5$;
30 wherein R^5 is selected from H,
 C_{1-6} -alkyl,
 C_{2-6} -alkenyl,
 C_{2-6} -alkynyl,
 C_{3-6} -cycloalkyl,

- 102 -

- C₄₋₆-cycloalkenyl,
R⁴,
phenyl optionally substituted with 1-3 substituents
independently selected from R³,
5 C₁₋₆-alkyl substituted with 1-3 substituents
independently selected from aryl, R⁷ and R⁴,
C₃₋₆-cycloalkyl substituted with 1-3 substituents
independently selected from aryl, R⁷ and R⁴, and
C₂₋₆-alkenyl substituted with 1-3 substituents
10 independently selected from aryl and R⁴;
wherein R⁶ is selected from C(O)R⁵, COOR⁵, C(O)NR⁵R⁵ and
S(O)_nR⁵;
wherein R⁷ is independently halo, OR⁵, NR⁵R⁵, COOR⁵ and CN.
- 15 3. The compound of claim 2 wherein R¹ is 5- or 6-
membered heteroaryl selected from isoxazolyl,
isothiazolyl, oxazolyl, pyranyl, thiazolyl, thiadiazolyl,
oxadiazolyl, furazanyl, furyl, and thienyl, wherein R¹ is
optionally substituted with 1-2 substituents
20 independently selected from R³;
wherein R² is selected from R⁴ and phenyl optionally
substituted with 1-4 substituents independently selected
from R³;
wherein R³ is independently selected from
25 C₁₋₄-alkyl, C₂₋₄-alkenyl, C₂₋₄-alkynyl, phenyl, R⁴, chloro,
fluoro, bromo, trifluoromethyl, C₁₋₄-alkoxy, phenoxy,
heterocyclyloxy, benzyloxy, C₁₋₄-alkylcarbonyloxy, amino,
alkylamino, phenylamino, carboxy, C₁₋₄-alkoxycarbonyl,
nitro, cyano, C₁₋₄-alkylcarbonyl, aminocarbonyl, C₁₋₄-
30 alkylaminocarbonyl, C₁₋₄-alkylsulfonyl, C₁₋₄-
alkylaminosulfonyl, benzyl, C₁₋₄-alkoxyalkyl, C₁₋₄-
aminoalkyl, C₁₋₄-alkylaminoalkyl, and 5-6- membered
heterocyclyl-C₁₋₄-alkyl;

- 103 -

wherein R⁴ is independently a 5-6 membered monocyclic, or 8-10 membered bicyclic saturated, partially saturated or unsaturated ring system comprising 1-3 heteroatoms if monocyclic, 1-6 heteroatoms if bicyclic, said heteroatoms independently selected from O, N, or S, which may be saturated or unsaturated, and wherein 0, 1, 2 or 3 atoms of each ring may be substituted by a substituent independently selected from C₁₋₄-alkyl, halo, C₁₋₆-haloalkyl, oxo, OR⁵, NR⁵R⁵, COOR⁵, nitro, cyano, S(O)_nR⁵, S(O)_nNR⁵R⁵, C(O)R⁵ and C(O)NR⁵R⁵; and wherein R⁵ is selected from H, C₁₋₆-alkyl, and phenyl optionally substituted with 1-3 substituents independently selected from C₁₋₄-alkyl, chloro, fluoro, -CF₃, hydroxy, C₁₋₄-alkoxy, amino, C₁₋₄-alkylamino, carboxy, C₁₋₄-alkoxycarbonyl, nitro, cyano, C₁₋₄-alkylcarbonyl, C₁₋₄-alkylaminocarbonyl, aminocarbonyl, aminosulfonyl and acetyl.

4. The compound of claim 3 wherein R¹ is selected from isoxazolyl, isothiazolyl, oxazolyl, thiazolyl, furyl, furazanyl and thienyl, wherein R¹ is optionally substituted with 1-2 substituents independently selected from R³;

wherein R² is selected from R⁴ and phenyl optionally substituted with 1-4 substituents independently selected from R³;

wherein R³ is independently selected from methyl, ethyl, propyl, tert-butyl, isopropyl, phenyl, chloro, fluoro, bromo, -CF₃, methoxy, phenoxy, benzyloxy, acetyl, amino, methylamino, phenylamino, carboxy, cyclohexyl, ethoxycarbonyl, nitro, cyano, methylcarbonyl, aminocarbonyl, methylaminocarbonyl, methylsulfonyl,


- 104 -

methylaminosulfonyl, benzyl, methoxymethyl, aminomethyl,
 N,N-dimethylaminoethyl and furylmethyl; and
 wherein R⁴ is independently selected from quinolyl,
 isoquinolyl, indazolyl, benzimidazolyl, pyrazolyl,
 5 pyrrolyl, indolyl, isoindolyl, purinyl, and
 naphthyridinyl, wherein R⁴ is optionally substituted by a
 substituent independently selected from methyl,
 isopropyl, tert-butyl, fluoro, chloro, -CF₃, oxo,
 methoxy, phenoxy, benzyloxy, amino, methylamino,
 10 phenylamino, carboxy, ethoxycarbonyl, nitro, cyano,
 methylcarbonyl, aminocarbonyl, methylaminocarbonyl,
 methylsulfonyl and methylaminosulfonyl.

5. Compound of Claim 4 wherein R² is selected from
 15 3,4,5-trimethoxyphenyl, 3,4-dimethoxyphenyl, 2,5-
 dimethoxyphenyl, 3,4-dimethoxy-6-methylphenyl, quinolinyl,
 benzimidazolyl, indazolyl, 3-aminosulfonylphenyl and 4-
 aminosulfonylphenyl.

20 6. Compound of Claim 4 wherein R¹ is isoxazolyl.

7. A compound of Formula II



II

25 wherein R¹⁰ is selected from phenyl, and 5-10 membered
 heterocyclyl; wherein R¹⁰ is optionally substituted with
 1-2 substituents selected from R¹¹;
 wherein R¹¹ is selected from C₁-C₄ alkyl, C₂-C₃ alkenyl, C₂-C₃
 alkynyl, C₃-C₆ cycloalkyl, C₄-C₆ cycloalkenyl, phenyl, 5-6

- 105 -

membered heterocyclyl, fluoro, chloro, CF_3 , $-\text{OR}^{12}$,
 $-\text{OC}(\text{O})\text{R}^{12}$, $-\text{NR}^{12}\text{R}^{12}$, $-\text{COOR}^2$, $-\text{C}(\text{O})\text{R}^{12}$, $-\text{C}(\text{O})\text{NR}^{12}\text{R}^{12}$, $-\text{SO}_2\text{R}^{12}$,
 $-\text{SO}_2\text{NR}^{12}\text{R}^{12}$, $-\text{NR}^{12}\text{C}(\text{O})\text{NR}^{12}\text{R}^{12}$, $-\text{NR}^{12}\text{C}(\text{O})\text{R}^2$, $-\text{NR}^{12}(\text{COOR}^{12})$,
 $-\text{NR}^{12}\text{SO}_2\text{NR}^{12}\text{R}^{12}$, $-\text{NR}^{12}\text{SO}_2\text{R}^{12}$, $-\text{OC}(\text{O})\text{NR}^{12}\text{R}^{12}$, $\text{C}_1\text{-C}_3$ alkyl

5 substituted with 1-3

substituents independently selected from optionally
substituted phenyl and optionally substituted 5-6
membered heterocyclyl; and

$\text{C}_2\text{-C}_3$ alkenyl substituted with 1-3 substituents

10 independently selected from optionally substituted
phenyl and optionally substituted 5-6 membered
heterocyclyl;

wherein R^{12} is selected from H, C_{1-6} -alkyl, and


phenyl optionally substituted with 1-3 substituents

15 independently selected from

C_{1-4} -alkyl, chloro, fluoro, $-\text{CF}_3$, hydroxy, C_{1-4} -
alkoxy, amino, C_{1-4} -alkylamino, carboxy, C_{1-4} -
alkoxycarbonyl, nitro, cyano, C_{1-4} -alkylcarbonyl,
 C_{1-4} -alkylaminocarbonyl, aminocarbonyl, aminosulfonyl
20 and acetyl;

and pharmaceutically acceptable derivative thereof.

8. A compound of Formula III



III

25 wherein R^{10} is selected from phenyl, and 5-10 membered
heterocyclyl; wherein R^{10} is optionally substituted with
1-2 substituents selected from R^{11} ;

wherein R^{11} is selected from $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_2\text{-C}_3$ alkenyl, $\text{C}_2\text{-C}_3$

30 alkynyl, $\text{C}_3\text{-C}_6$ cycloalkyl, $\text{C}_4\text{-C}_6$ cycloalkenyl, phenyl, 5-6

- 106 -

membered heterocyclyl, fluoro, chloro, CF_3 , $-\text{OR}^{12}$,
 $-\text{OC}(\text{O})\text{R}^{12}$, $-\text{NR}^{12}\text{R}^{12}$, $-\text{COOR}^2$, $-\text{C}(\text{O})\text{R}^{12}$, $-\text{C}(\text{O})\text{NR}^{12}\text{R}^{12}$, $-\text{SO}_2\text{R}^{12}$,
 $-\text{SO}_2\text{NR}^{12}\text{R}^{12}$, $-\text{NR}^{12}\text{C}(\text{O})\text{NR}^{12}\text{R}^{12}$, $-\text{NR}^{12}\text{C}(\text{O})\text{R}^2$, $-\text{NR}^{12}(\text{COOR}^{12})$,
 $-\text{NR}^{12}\text{SO}_2\text{NR}^{12}\text{R}^{12}$, $-\text{NR}^{12}\text{SO}_2\text{R}^{12}$, $-\text{OC}(\text{O})\text{NR}^{12}\text{R}^{12}$, C_1 - C_3 alkyl

5 substituted with 1-3

substituents independently selected from optionally
substituted phenyl and optionally substituted 5-6
membered heterocyclyl; and

C_2 - C_3 alkenyl substituted with 1-3 substituents

10 independently selected from optionally substituted
phenyl and optionally substituted 5-6 membered
heterocyclyl;

wherein R^{12} is selected from H, C_{1-6} -alkyl, and


phenyl optionally substituted with 1-3 substituents

15 independently selected from

C_{1-4} -alkyl, chloro, fluoro, CF_3 , hydroxy, C_{1-4} -alkoxy,
amino, C_{1-4} -alkylamino, carboxy, C_{1-4} -alkoxycarbonyl,
nitro, cyano, C_{1-4} -alkylcarbonyl, C_{1-4} -
alkylaminocarbonyl, aminocarbonyl, aminosulfonyl and
20 acetyl;

and pharmaceutically acceptable derivatives thereof.

9. A compound of Formula I'



wherein R^1 is heteroaryl containing at least one oxygen or
sulfur heteroatom, wherein R^1 is optionally substituted
with 1-4 substituents

30 wherein R^2 is selected from R^4 and aryl optionally
substituted with 1-5 substituents independently selected
from R^3 ;

- 107 -

wherein R^3 is independently selected from

H,
C₁₋₁₀-alkyl,
C₂₋₁₀-alkenyl,
5 C₂₋₁₀-alkynyl,
C₁₋₁₀-haloalkyl,
C₃₋₁₀-cycloalkyl,
C₄₋₁₀-cycloalkenyl,
aryl,
10 R^4 ,
halo,
 SR^5 ,
 OR^5 ,
 $OC(O)R^5$,
15 NR^5R^5 ,
 NR^5R^6 ,
 $COOR^5$,
nitro,
cyano,
20 $C(O)R^5$,
 $C(O)C(O)R^5$,
 $C(O)NR^5R^5$,
 $S(O)_nR^5$,
 $S(O)_nNR^5R^5$,
25 $S(O)_nNR^5R^6$,
 $NR^5C(O)NR^5R^5$,
 $NR^5C(O)C(O)R^5$,
 $NR^5C(O)R^5$,
 $NR^5(COOR^5)$,
30 $NR^5C(O)R^4$,
 $NR^5S(O)_nNR^5R^5$,
 $NR^5S(O)_nR^5$,
 $NR^5S(O)_nR^4$,
 $NR^5C(O)C(O)NR^5R^5$,

- 108 -

$\text{NR}^5\text{C}(\text{O})\text{C}(\text{O})\text{NR}^5\text{R}^6$,
 C_{1-10} -alkyl substituted with 1-3 substituents
independently selected from aryl, R^7 and R^4 ; and
 C_{2-10} -alkenyl substituted with 1-3 substituents
5 independently selected from aryl, R^7 and R^4 ;
wherein R^4 is independently a 5-8 membered monocyclic, 8-12
membered bicyclic, or 11-14 membered tricyclic saturated,
partially saturated or unsaturated ring system comprising
1-3 heteroatoms if monocyclic, 1-6 heteroatoms if
10 bicyclic, or 1-9 heteroatoms if tricyclic, said
heteroatoms independently selected from O, N, or S, which
may be saturated or unsaturated, and wherein 0, 1, 2 or 3
atoms of each ring may be substituted by a substituent
independently selected from C_{1-10} -alkyl, C_{2-10} -alkenyl, C_{2-}
15 $_{10}$ -alkynyl, C_{3-10} -cycloalkyl, C_{4-10} -cycloalkenyl, halo,
haloalkyl, sulfo, oxo, SR^5 , OR^5 , NR^5R^5 , NR^5R^6 , NR^6R^6 , COOR^5 ,
nitro, cyano, $\text{S}(\text{O})_n\text{R}^5$, $\text{S}(\text{O})_n\text{NR}^5\text{R}^5$; $\text{C}(\text{O})\text{R}^5$, $\text{C}(\text{O})\text{NR}^5\text{R}^5$ and 6-
membered heteroaryl optionally substituted with 1-3
substituents independently selected from R^3 ;
20 wherein R^5 is independently selected from H, C_{1-10} -alkyl, C_{2-}
 $_{10}$ -alkenyl, C_{2-10} -alkynyl, C_{3-10} -cycloalkyl, C_{4-10} -
cycloalkenyl, R^4 , aryl optionally substituted
with 1-3 substituents independently selected from R^3 ,
 $\text{C}_1\text{-C}_{10}$ alkyl substituted with 1-3 substituents
25 independently selected from aryl, R^7 and R^4 ;
 $\text{C}_3\text{-C}_{10}$ cycloalkyl substituted with 1-3 substituents
independently selected from aryl, R^7 and R^4 ; and
 $\text{C}_2\text{-C}_{10}$ alkenyl substituted with 1-3 substituents
independently selected from aryl, R^7 and R^4 ;
30 wherein R^6 is selected from $-\text{C}(\text{O})\text{R}^5$, $-\text{COOR}^5$, $-\text{C}(\text{O})\text{NR}^5\text{R}^5$ and
 $-\text{S}(\text{O})_n\text{R}^5$;
wherein R^7 is independently halo, $-\text{CF}_3$, $-\text{SR}^5$, $-\text{OR}^5$, $-\text{OC}(\text{O})\text{R}^5$,
 $-\text{NR}^5\text{R}^5$, $-\text{NR}^5\text{R}^6$, $-\text{NR}^6\text{R}^6$, $-\text{COOR}^5$, $-\text{NO}_2$, $-\text{CN}$, $-\text{C}(\text{O})\text{R}^5$,

- 109 -

$-\text{OC}(\text{O})\text{NR}^5\text{R}^5$, $-\text{C}(\text{O})\text{NR}^5\text{R}^5$, $-\text{N}(\text{R}^5)\text{C}(\text{O})\text{R}^5$, $-\text{N}(\text{R}^5)(\text{COOR}^5)$ and $-\text{S}(\text{O})_n\text{NR}^5\text{R}^5$; and wherein n is 1 or 2;

and pharmaceutically acceptable derivative thereof; provided both R^1 and R^2 are not 5-*t*-butyl-3-isoxazolyl.

5

10. The compound of claim 9, wherein R^1 is 5- or 6-membered heteroaryl containing at least one oxygen or sulfur heteroatom, wherein R^1 is optionally substituted with 1-3 substituents independently selected from R^3 ;
- 10 wherein R^2 is selected from R^4 and aryl optionally substituted with 1-3 substituents independently selected from R^3 ;
- wherein R^3 is independently selected from C_{1-4} -alkyl, C_{2-4} -alkenyl, C_{2-4} -alkynyl, phenyl, C_{6-10} -cycloalkyl, R^4 , chloro,
- 15 fluoro, bromo, trifluoromethyl, C_{1-4} -alkoxy, phenoxy, heterocyclyloxy, benzyloxy, C_{1-4} -alkylcarbonyloxy, amino, alkylamino, phenylamino, carboxy, C_{1-4} -alkoxycarbonyl, nitro, cyano, C_{1-4} -alkylcarbonyl, aminocarbonyl, C_{1-4} -alkylaminocarbonyl, C_{1-4} -alkylsulfonyl, C_{1-4} -
- 20 alkylaminosulfonyl, benzyl, C_{1-4} -alkoxyalkyl, C_{1-4} -aminoalkyl, C_{1-4} -alkylaminoalkyl, and 5-6- membered heterocyclyl- C_{1-4} -alkyl;
- wherein R^4 is independently 2,3-dihydro-indolyl, 1,3-benzodioxolyl, indolyl, 1,3-dioxo-isoindolyl, indazolyl,
- 25 pyridyl, quinolyl, isoquinolyl, benzothiazolyl, 1,2,3-benzotriazolyl, benzimidazolyl, and pyridyl; wherein R^4 is optionally substituted with hydroxy, C_{1-3} -alkoxy, cyano, nitro, halo, C_{1-3} -alkyl, di- C_{1-3} -alkylamino, di- C_{1-3} -alkylamino- C_{1-3} -alkyl, di- C_{1-3} -alkylamino- C_{1-3} -alkoxy, C_{1-3} -
- 30 alkylcarbonyl, C_{1-3} -alkoxycarbonyl, C_{1-3} -alkylcarbonylamino, pyrrolidinylcarbonyl- C_{2-3} -alkenyl, pyrrolidinylcarbonyl- C_{1-3} -alkyl, pyrrolidinyl- C_{1-3} -alkyl, C_{2-3} -alkynyl, acetyl, C_{1-3} -alkylcarbonyl- C_{1-3} -alkyl, carboxy- C_{1-3} -alkyl, (piperidinyl)- C_{1-3} -alkoxy,

- 110 -

(piperazinyl)-C₁₋₃-alkoxy, 2-morpholinyl-C₁₋₃-alkoxy, C₁₋₃-haloalkyl, C₁₋₃-haloalkoxy, aminocarbonyl, aminosulfonyl, C₁₋₃-alkylaminosulfonyl, hydroxy-C₁₋₃-alkylaminosulfonyl, (thiazolyl)aminosulfonyl, C₁₋₄-alkylaminosulfonyl, C₁₋₃-alkylcarbonylaminosulfonyl, C₁₋₃-alkylsulfonyl, C₁₋₃-alkoxycarbonyl-piperazinyl, morpholinyl, C₁₋₃-alkylpiperziny, piperazinyl, C₁₋₃-alkyl-piperazinyl, and oxazolyl.

10 11. The compound of claim 10 wherein R¹ is selected from isoxazolyl, isothiazolyl, oxazolyl, pyranyl, thiazolyl, thiadiazolyl, oxadiazolyl, furazanyl, furyl, and thienyl, wherein R¹ is optionally substituted with 1-2 substituents independently selected from R³;
15 wherein R² is selected from naphthyl, 2,3-dihydro-indolyl, 1,3-benzodioxolyl, indolyl, 1,3-dioxo-isindolyl, indazolyl, pyridyl, quinolyl, isoquinolyl, benzothiazolyl, 1,2,3-benzotriazolyl, benzimidazolyl, and phenyl; wherein R² is optionally substituted with 1-3
20 substituents independently selected from R³;
 wherein R³ is independently selected from H, halo, C₁₋₃-alkyl, C₂₋₃-alkenyl, C₂₋₃-alkynyl, phenyl, C₆₋₁₀-cycloalkyl, hydroxy, C₁₋₃-haloalkoxy, C₁₋₃-alkoxy, -C(O)-C₁₋₃-alkyl, and C₁₋₃-haloalkyl.

25 12. The compound of claim 11 wherein R¹ is selected from isoxazolyl, isothiazolyl, oxazolyl, thiazolyl, furyl, furazanyl and thienyl, wherein R¹ is optionally substituted with 1-2 substituents independently selected
30 from R³;
 wherein R² is selected from 2-naphthyl, 2,3-dihydro-indol-6-yl, 1,3-benzodioxol-5-yl, 5-indolyl, 4-indolyl, 1,3-dioxo-isindol-5-yl, 5-indazolyl, 6-indazolyl, 3-pyridyl, 3-quinolyl, 6-quinolyl, isoquinolyl, benzothiazol-6-yl,

- 111 -

benzothiazol-5-yl, 1,2,3-benzotriazol-5-yl, 6-benzimidazolyl, 5-pyridyl, and phenyl;

wherein R² is optionally substituted with 1-3 substituents

independently selected from hydroxy, methoxy, ethoxy,
5 cyano, nitro, chloro, fluoro, bromo, dimethylamino, dimethylaminoethyl, 3-dimethylamino-propoxy, methoxycarbonyl, methylcarbonyl, methylcarbonylamino, CH₃C(O)N(CH₃), methyl, ethyl, isopropyl, pyrrolidin-1-ylcarbonylethenyl, pyrrolidin-1-ylcarbonylethyl,
10 pyrrolidin-1-ylpropyl, ethynyl, acetyl, ethoxycarbonylbutyl, carboxybutyl, 2-(1-methyl-piperidin-4-yl)-ethoxy, 2-(4-methyl-piperazin-1-yl)ethoxy, 3-(4-methyl-piperazin-1-yl)-propoxy, 3-(piperidin-1-yl)propoxy, 2-piperidin-1-yl-ethoxy, 2-morpholin-4-yl-ethoxy, pentafluoroethyl, trifluoromethyl,
15 trifluoromethoxy, difluoromethoxy, aminocarbonyl, aminosulfonyl, N,N'-di-propylaminosulfonyl, hydroxypropylaminosulfonyl, (2-thiazolyl)aminosulfonyl, butylaminosulfonyl, methylcarbonylaminosulfonyl, methylsulfonyl, 1-methyl-piperidin-4-ylmethoxy, 1-tert-butoxycarbonyl-piperazin-4-yl, 4-morpholinyl, 4-methylpiperzine-1-yl, 4-piperazinyl, 4-isopropyl-piperazin-1-yl, and oxazol-5-yl; and
20 wherein R³ is independently selected from H, hydroxy, iodo, methyl, acetyl, trifluoromethyl, methoxy, cyclohexyl, adamantyl, phenyl and trifluoromethoxy.

13. Compound of Claim 9 wherein R² is selected from 3,4,5-trimethoxyphenyl, 3,4-dimethoxyphenyl, 2,5-dimethoxyphenyl, 3,4-dimethoxy-6-methylphenyl, 3-methylcarbonylamino-phenyl, 4-[CH₃C(O)NH-SO₂]-phenyl, 3-isopropylphenyl, 4-[CH₃C(O)N(CH₃)]phenyl, 3-methoxy-4-morpholinylphenyl, 3-quinolinyl, 6-quinolinyl,
30

- 112 -

benzimidazolyl, 5-indazolyl, 6-indazolyl, 3-chlorophenyl, 3-aminosulfonylphenyl and 4-aminosulfonylphenyl.

14. Compound of Claim 9 wherein R¹ is substituted
5 isoxazolyl.

15 15. Compound of Claim 9 wherein R¹ is 3-isoxazolyl substituted with a substituent selected from cyclohexyl, adamantyl, phenyl and tert-butyl.

16. Compound of Claim 9 wherein R¹ is 5-isoxazolyl substituted with a substituent selected from cyclohexyl, adamantyl, phenyl and tert-butyl.


15 17. Compound of Claim 9 and pharmaceutically acceptable derivatives thereof selected from

N⁴-(5-tert-butylisoxazol-3-yl)-N²-(3,4,5-trimethoxyphenyl)-pyrimidine-2,4-diamine;
20 N⁴-(5-t-butyl-isoxazol-3-yl)-N²-(4,5-dimethoxy-2-methylphenyl)-pyrimidine-2,4-diamine;
N⁴-(5-t-butyl-isoxazol-3-yl)-N²-(1H-indazol-5-yl)-pyrimidine-2,4-diamine;
N²-(3-methoxy-4-morpholin-4-yl-phenyl)-N⁴-(5-phenyl-
25 isoxazol-3-yl)-pyrimidine-2,4-diamine;
N²-(3,4,5-trimethoxyphenyl)-N⁴-(5-phenylisoxazol-3-yl)-2,4-pyrimidinediamine;
N²-(2-methyl-4,5-dimethoxyphenyl)-N⁴-(5-phenylisoxazol-3-yl)-2,4-pyrimidinediamine;
30 N²-(4,5-dimethoxy-2-methylphenyl)-N⁴-(5-adamantylisoxazol-3-yl)-2,4-pyrimidinediamine; and
N²-(3,4,5-trimethoxyphenyl)-N⁴-(3-cyclohexylisoxazol-5-yl)-2,4-pyrimidinediamine.

35

- 113 -

18. A compound of Formula II'



II'


- 5 wherein R^{10} is selected from phenyl, and 5-10 membered heterocyclyl; wherein R^{10} is optionally substituted with 1-2 substituents selected from R^{11} ;
- wherein R^{11} is selected from C_1 - C_4 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_3 - C_{10} cycloalkyl, C_4 - C_6 cycloalkenyl, phenyl, 5-6
- 10 membered heterocyclyl, fluoro, chloro, CF_3 , $-OC(O)R^{12}$, $-NR^{12}R^{12}$, $-COOR^2$, $-C(O)R^{12}$, $-C(O)NR^{12}R^{12}$, $-SO_2R^{12}$, $-SO_2NR^{12}R^{12}$, $-NR^{12}C(O)NR^{12}R^{12}$, $-NR^{12}C(O)R^2$, $-NR^{12}(COOR^{12})$, $-NR^{12}SO_2NR^{12}R^{12}$, $-NR^{12}SO_2R^{12}$, $-OC(O)NR^{12}R^{12}$, $-OR^{13}$, C_1 - C_3 alkyl substituted with 1-3 substituents independently selected
- 15 from optionally substituted phenyl and optionally substituted 5-6 membered heterocyclyl; and C_2 - C_3 alkenyl substituted with 1-3 substituents independently selected from optionally substituted phenyl and optionally substituted 5-6 membered
- 20 heterocyclyl;
- wherein R^{12} is selected from H, C_1 - C_6 -alkyl, and phenyl optionally substituted with 1-3 substituents independently selected from
- 25 C_1 - C_4 -alkyl, chloro, fluoro, CF_3 , hydroxy, C_1 - C_4 -alkoxy, amino, C_1 - C_4 -alkylamino, carboxy, C_1 - C_4 -alkoxycarbonyl, NO_2 , CN, C_1 - C_4 -alkylcarbonyl, C_1 - C_4 -alkylaminocarbonyl, aminocarbonyl, aminosulfonyl and acetyl; and

- 114 -

wherein R^{13} is selected from H, C_{1-6} -alkyl, amino- C_{1-6} -alkyl, C_{1-6} -alkylamino- C_{1-6} -alkyl, aminocarbonyl- C_{1-6} -alkyl, C_{1-6} -alkylaminocarbonyl- C_{1-6} -alkyl, and phenyl optionally substituted with 1-3 substituents independently
 5 selected from C_{1-4} -alkyl, chloro, fluoro, CF_3 , hydroxy, C_{1-4} -alkoxy, amino, C_{1-4} -alkylamino, carboxy, C_{1-4} -alkoxycarbonyl, NO_2 , CN, C_{1-4} -alkylcarbonyl, C_{1-4} -alkylaminocarbonyl, aminocarbonyl, aminosulfonyl and acetyl; and pharmaceutically acceptable salts
 10 thereof; provided R^{10} is not 5-*t*-butyl-3-isoxazolyl when R^{11} is *t*-butyl.

19. Compound of Claim 18 wherein R^{10} is selected from 3,4,5-trimethoxyphenyl, 3,4-dimethoxy-6-methylphenyl, 3-
 15 methoxy-4-morpholinylphenyl and 5-indazolyl; and wherein R^{11} is selected from cyclohexyl, adamantyl, phenyl and *tert*-butyl.

20. A compound of Formula III'



III'

wherein R^{10} is selected from phenyl, and 5-10 membered heterocyclyl; wherein R^{10} is optionally substituted with
 25 1-2 substituents selected from R^{11} ;
 wherein R^{11} is selected from C_1 - C_4 alkyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, C_3 - C_{10} cycloalkyl, C_4 - C_6 cycloalkenyl, phenyl, 5-6 membered heterocyclyl, fluoro, chloro, CF_3 , $-OC(O)R^{12}$, $-NR^{12}R^{12}$, $-COOR^2$, $-C(O)R^{12}$, $-C(O)NR^{12}R^{12}$, $-SO_2R^{12}$, $-SO_2NR^{12}R^{12}$,
 30 $-NR^{12}C(O)NR^{12}R^{12}$, $-NR^{12}C(O)R^2$, $-NR^{12}(COOR^{12})$, $-NR^{12}SO_2NR^{12}R^{12}$,

- 115 -

-NR¹²SO₂R¹², -OC(O)NR¹²R¹², -OR¹³, C₁-C₃ alkyl substituted with 1-3 substituents independently selected from optionally substituted phenyl and optionally substituted 5-6 membered heterocyclyl; and

5 C₂-C₃ alkenyl substituted with 1-3 substituents independently selected from optionally substituted phenyl and optionally substituted 5-6 membered heterocyclyl;

wherein R¹² is selected from H, C₁₋₆-alkyl, and

10 phenyl optionally substituted with 1-3 substituents independently selected from C₁₋₄-alkyl, chloro, fluoro, CF₃, hydroxy, C₁₋₄-alkoxy, amino, C₁₋₄-alkylamino, carboxy, C₁₋₄-alkoxycarbonyl, NO₂, CN, C₁₋₄-alkylcarbonyl, C₁₋₄-alkylaminocarbonyl,

15 aminocarbonyl, aminosulfonyl and acetyl; and

wherein R¹³ is selected from H, C₁₋₆-alkyl, amino-C₁₋₆-alkyl, C₁₋₆-alkylamino-C₁₋₆-alkyl, aminocarbonyl-C₁₋₆-alkyl, C₁₋₆-alkylaminocarbonyl-C₁₋₆-alkyl, and phenyl optionally substituted with 1-3 substituents independently

20 selected from C₁₋₄-alkyl, chloro, fluoro, CF₃, hydroxy, C₁₋₄-alkoxy, amino, C₁₋₄-alkylamino, carboxy, C₁₋₄-alkoxycarbonyl, NO₂, CN, C₁₋₄-alkylcarbonyl, C₁₋₄-alkylaminocarbonyl, aminocarbonyl, aminosulfonyl and acetyl; and


25 pharmaceutically acceptable salts thereof; provided R¹⁰ is not 5-*t*-butyl-3-isoxazolyl when R¹¹ is *t*-butyl.

21. Compound of Claim 20 wherein R¹⁰ is selected from

30 3,4,5-trimethoxyphenyl, 3,4-dimethoxy-6-methylphenyl, 3-methoxy-4-morpholinylphenyl and 5-indazolyl; and wherein R¹¹ is selected from cyclohexyl, adamantyl, phenyl and *tert*-butyl.

- 116 -

22. A compound of Formula IV



IV

- 5 wherein R^{10} is selected from phenyl, and 5-10 membered heterocyclyl; wherein R^{10} is optionally substituted with 1-2 substituents selected from R^{11} ;
- wherein R^{11} is selected from $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_2\text{-C}_3$ alkenyl, $\text{C}_2\text{-C}_3$ alkynyl, $\text{C}_3\text{-C}_{10}$ cycloalkyl, $\text{C}_4\text{-C}_6$ cycloalkenyl, phenyl, 5-6
- 10 membered heterocyclyl, fluoro, chloro, CF_3 , $-\text{OC}(\text{O})\text{R}^{12}$, $-\text{NR}^{12}\text{R}^{12}$, $-\text{COOR}^2$, $-\text{C}(\text{O})\text{R}^{12}$, $-\text{C}(\text{O})\text{NR}^{12}\text{R}^{12}$, $-\text{SO}_2\text{R}^{12}$, $-\text{SO}_2\text{NR}^{12}\text{R}^{12}$, $-\text{NR}^{12}\text{C}(\text{O})\text{NR}^{12}\text{R}^{12}$, $-\text{NR}^{12}\text{C}(\text{O})\text{R}^2$, $-\text{NR}^{12}(\text{COOR}^{12})$, $-\text{NR}^{12}\text{SO}_2\text{NR}^{12}\text{R}^{12}$, $-\text{NR}^{12}\text{SO}_2\text{R}^{12}$, $-\text{OC}(\text{O})\text{NR}^{12}\text{R}^{12}$, $-\text{OR}^{13}$, $\text{C}_1\text{-C}_3$ alkyl substituted with 1-3 substituents independently selected
- 15 from optionally substituted phenyl and optionally substituted 5-6 membered heterocyclyl; and
- $\text{C}_2\text{-C}_3$ alkenyl substituted with 1-3 substituents independently selected from optionally substituted phenyl and optionally substituted 5-6 membered
- 20 heterocyclyl;
- wherein R^{12} is selected from H, C_{1-6} -alkyl, and phenyl optionally substituted with 1-3 substituents independently selected from
- C_{1-4} -alkyl, chloro, fluoro, CF_3 , hydroxy, C_{1-4} -alkoxy,
- 25 amino, C_{1-4} -alkylamino, carboxy, C_{1-4} -alkoxycarbonyl, NO_2 , CN, C_{1-4} -alkylcarbonyl, C_{1-4} -alkylaminocarbonyl, aminocarbonyl, aminosulfonyl and acetyl; and
- wherein R^{13} is selected from H, C_{1-6} -alkyl, amino- C_{1-6} -alkyl, C_{1-6} -alkylamino- C_{1-6} -alkyl, aminocarbonyl- C_{1-6} -alkyl, C_{1-6} -alkylaminocarbonyl- C_{1-6} -alkyl, and phenyl optionally
- 30

- 117 -

substituted with 1-3 substituents independently
selected from

C₁₋₄-alkyl, chloro, fluoro, CF₃, hydroxy, C₁₋₄-alkoxy,
amino, C₁₋₄-alkylamino, carboxy, C₁₋₄-alkoxycarbonyl,
5 NO₂, CN, C₁₋₄-alkylcarbonyl, C₁₋₄-alkylaminocarbonyl,
aminocarbonyl, aminosulfonyl and acetyl; and
pharmaceutically acceptable salts thereof.

23. Compound of Claim 22 wherein R¹⁰ is selected from
10 3,4,5-trimethoxyphenyl, 3,4-dimethoxy-6-methylphenyl, 3-
methoxy-4-morpholinylphenyl and 5-indazolyl; and wherein R¹¹
is selected from cyclohexyl, adamantyl, phenyl and tert-
butyl.

15 24. A composition comprising a compound according to
any of claims 1-23 and a pharmaceutically acceptable
carrier.

25. A method of treating IGF-1R related disorders in a
20 mammal, said method comprising administering an effective
amount of a compound of any of Claims 1-23.

26. A method of treating a kinase mediated disease or
disease symptoms in a mammal comprising administration to
25 said mammal of a compound according to any of claims 1-23.

27. A method of inhibiting kinase activity in a
mammal comprising the step of administering to said mammal a
compound according to any of claims 1-23.

30

28. A method of treating cancer in a subject, said
method comprising administering an effective amount of a
compound of any of Claims 1-23.

- 118 -

29. A compound as in any of Claims 1-23 for use in a method of therapeutic treatment for the human or animal body.

5 30. A method of reducing tumor size in a subject, said method comprising administering an effective amount of a compound as in any of Claims 1-23.

10 31. Use of a compound of any of Claims 1-23 for preparing a medicament for the treatment of cancer.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/26926

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/506 A61P35/00 C07D413/12 C07D413/14 C07D417/14
C07D417/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 02 45652 A (BILODEAU MARK T ;MANLEY PETER J (US); MERCK & CO INC (US); RODMAN) 13 June 2002 (2002-06-13) Formula I, ex. 2-3	1,7-9, 18,20, 22,24, 26-31
P,X	WO 01 64655 A (BREAULT GLORIA ANNE ;PEASE ELIZABETH JANET (GB); ASTRAZENECA UK LT) 7 September 2001 (2001-09-07) Formula (I)	1,7-9, 18,20, 22,24, 26-31
X	US 5 958 935 A (DAVIS JEREMY MARTIN ET AL) 28 September 1999 (1999-09-28) cited in the application Formula (I)	1,7-9, 18,20, 22,24, 26-31



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

*" document member of the same patent family

Date of the actual completion of the international search

13 November 2002

Date of mailing of the international search report

25/11/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Johnson, C

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 02/26926

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 01 40218 A (AVENTIS PHARMA SA) 7 June 2001 (2001-06-07) cited in the application claim 1 ---	1,7-9, 18,20, 22,24, 26-31
A	WO 00 39101 A (BREault GLORIA ANNE ;PEASE JANET ELIZABETH (GB); ASTRAZENECA UK LT) 6 July 2000 (2000-07-06) cited in the application Formula (I), examples 134 and 159 -----	1,7-9, 18-20, 22,24, 26-31

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claims 25-28, 30 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition. Although claim(s) are directed to a diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box I.2

Claims Nos.: 1(part),7(part),8(part),9(part),18(part),20(part)22(part)

The term "pharmaceutically acceptable derivatives thereof" used extensively throughout the claims is not clear and hence does not fulfil the requirements of Article 6 PCT. The definition on p. 28 of the description simply provides a definition in functional terms, i.e. a compound that is capable of providing, directly or indirectly, a compound of the invention (or a metabolite or residue thereof), but does not provide any information concerning the structure of such a derivative. In view of the lack of clarity, the search has been restricted to compounds as defined by the given formulae and their pharmaceutically acceptable salts.

It should be noted that the application does not fulfil the requirements of Article 6 PCT (clarity and conciseness) as there are 7 compound claims which are presented as being independent. Some of these claims appear in fact to be dependent claims, whilst others overlap with one another. In view of the confused method of claiming, the search has concentrated on those claims presented as being independent (i.e. the further features of the dependent claims have not been searched).

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 02/26926

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☒ Claims Nos.: 1(part), 7(part), 8(part), 9(part), 18(part), 20(part) 22(part)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 02/26926

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0245652	A	13-06-2002	AU 3244102 A	18-06-2002
			WO 0245652 A2	13-06-2002
			US 2002137755 A1	26-09-2002
WO 0164655	A	07-09-2001	AU 3397601 A	12-09-2001
			WO 0164655 A1	07-09-2001
US 5958935	A	28-09-1999	AU 7631496 A	11-06-1997
			EP 0862560 A1	09-09-1998
			WO 9719065 A1	29-05-1997
			US 6235746 B1	22-05-2001
WO 0140218	A	07-06-2001	FR 2801588 A1	01-06-2001
			AU 2179001 A	12-06-2001
			BR 0015992 A	06-08-2002
			CZ 20021849 A3	14-08-2002
			EP 1244650 A1	02-10-2002
			WO 0140218 A1	07-06-2001
			NO 20022528 A	28-05-2002
WO 0039101	A	06-07-2000	AU 1874300 A	31-07-2000
			BR 9916590 A	23-10-2001
			CN 1335838 T	13-02-2002
			EP 1140860 A1	10-10-2001
			WO 0039101 A1	06-07-2000
			JP 2002533446 T	08-10-2002
			NO 20013038 A	22-08-2001

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
6 March 2003 (06.03.2003)

PCT

(10) International Publication Number
WO 03/018022 A1(51) International Patent Classification⁷: A61K 31/506,
A61P 35/00, C07D 413/14, 413/12, 417/14, 417/12

(21) International Application Number: PCT/US02/26926

(22) International Filing Date: 22 August 2002 (22.08.2002)

(25) Filing Language: English

(26) Publication Language: English


(30) Priority Data:
60/314,285 22 August 2001 (22.08.2001) US
10/225,819 21 August 2002 (21.08.2002) US(71) Applicant: AMGEN INC. [US/US]; One Amgen Center
Drive, Thousand Oaks, CA 91320-1799 (US).(72) Inventors: BUCHANAN, John, L.; 143 Babcock Street,
Brookline, MA 02446 (US). CHAFFEE, Stuart; 24 Bates
Street, Cambridge, MA 02140 (US). HARMANGE, Jean-
Christophe; 57 William Street, Andover, MA 01810 (US).
NOVAK, Perry, M.; 35 Debbie Lane, Milford, MA 01757
(US). ZHU, Xiaotian; 131 Coolidge Avenue #324, Water-
town, MA 02472 (US).(74) Agents: ODRE, Steven, M. et al.; Amgen, Inc., One Am-
gen Center Drive, M/S 27-4-A, Thousand Oaks, CA 91320-
1799 (US).(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN,
YU, ZA, ZM, ZW.(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(48) Date of publication of this corrected version:
2 October 2003(15) Information about Correction:
see PCT Gazette No. 40/2003 of 2 October 2003, Section IIFor two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: 2-AMINO-4-HETEROARYLAMINOPYRIMIDINE DERIVATIVES FOR USE IN THE TREATMENT OF CANCER



(I)

(57) Abstract: Compounds of Formula (I) are claimed, wherein R¹ is
a heteroaryl ring containing at least one oxygen or sulfur heteroatom
and R² is as defined in the claims. These compounds are useful in the
treatment of cancer.

WO 03/018022 A1